

QA 343.B16 C.1

Elliptic func

00300



3 1735 015 185 105

ELLIPTIC FUNCTIONS.

AN ELEMENTARY TEXT-BOOK FOR
STUDENTS OF MATHEMATICS.

BY

ARTHUR L. BAKER, C.E., PH.D.,

HEAD OF DEPARTMENT OF MATHEMATICS, MANUAL TRAINING HIGH SCHOOL, BROOKLYN, N. Y.;
FORMERLY PROFESSOR OF MATHEMATICS IN THE UNIVERSITY OF ROCHESTER, N. Y.

$$\sin \operatorname{am} u = \frac{1}{\sqrt{k}} \cdot \frac{H(u)}{\Theta(u)}.$$

FIRST EDITION.

FIRST THOUSAND.

NEW YORK:

JOHN WILEY & SONS.

LONDON: CHAPMAN & HALL, LIMITED.

1906

Copyright, 1890,
BY
ARTHUR L. BAKER.

ROBERT DRUMMOND, PRINTER, NEW YORK

PREFACE.

IN the works of Abel, Euler, Jacobi, Legendre, and others, the student of Mathematics has a most abundant supply of material for the study of the subject of Elliptic Functions.

These works, however, are not accessible to the general student, and, in addition to being very technical in their treatment of the subject, are moreover in a foreign language.

It is in the hope of smoothing the road to this interesting and increasingly important branch of Mathematics, and of putting within reach of the English student a tolerably complete outline of the subject, clothed in simple mathematical language and methods, that the present work has been compiled.

New or original methods of treatment are not to be looked for. The most that can be expected will be the simplifying of methods and the reduction of them to such as will be intelligible to the average student of Higher Mathematics.

I have endeavored throughout to use only such methods as are familiar to the ordinary student of Calculus, avoiding those methods of discussion dependent upon the properties of double periodicity, and also those depending upon Functions of Complex Variables. For the same reason I have not carried the discussion of the Θ and H functions further.

Among the minor helps to simplicity is the use of zero subscripts to indicate decreasing series in the Landen Transformation, and of numerical subscripts to indicate increasing series. I have adopted the notation of Gudermann, as being more simple than that of Jacobi.

I have made free use of the following works: JACOBI'S *Fundamenta Nova Theoriæ Func. Ellip.*; HOUEL'S *Calcul Infinitésimal*; LEGENDRE'S *Traité des Fonctions Elliptiques*; DUREGE'S *Theorie der Elliptischen Functionen*; HERMITE'S *Théorie des Fonctions Elliptiques*; VERHULST'S *Théorie des Fonctions Elliptiques*; BERTRAND'S *Calcul Intégral*; LAURENT'S *Théorie des Fonctions Elliptiques*; CAYLEY'S *Elliptic Functions*; BYERLY'S *Integral Calculus*; SCHLOMILCH'S *Die Höheren Analysis*; BRIOT ET BOUQUET'S *Fonctions Elliptiques*.

I have refrained from any reference to the Gudermann or Weierstrass functions as not within the scope of this work, though the Gudermannians might have been interesting examples of verification formulæ. The arithmetico-geometrical mean, the march of the functions, and other interesting investigations have been left out for want of room.

CONTENTS.

INTRODUCTORY CHAPTER,	o
CHAP. I. ELLIPTIC INTEGRALS,	o
II. ELLIPTIC FUNCTIONS,	o
III. PERIODICITY OF THE FUNCTIONS,	o
IV. LANDEN'S TRANSFORMATION,	o
V. COMPLETE FUNCTIONS,	o
VI. EVALUATION FOR ϕ ,	o
VII. FACTORIZATION OF ELLIPTIC FUNCTIONS,	o
VIII. THE Θ FUNCTION,	o
IX. THE Θ AND H FUNCTIONS,	o
X. ELLIPTIC INTEGRALS OF THE SECOND ORDER,	o
XI. ELLIPTIC INTEGRALS OF THE THIRD ORDER,	o
XII. NUMERICAL CALCULATIONS, q ,	o
XIII. NUMERICAL CALCULATIONS, K ,	o
XIV. NUMERICAL CALCULATIONS, u ,	o
XV. NUMERICAL CALCULATIONS, ϕ ,	o
XVI. NUMERICAL CALCULATIONS, $E(k, \phi)$,	o
XVII. APPLICATIONS,	o

ELLIPTIC FUNCTIONS.

INTRODUCTORY CHAPTER.*

THE first step taken in the theory of Elliptic Functions was the determination of a relation between the amplitudes of three functions of either order, such that there should exist an algebraic relation between the three functions themselves of which these were the amplitudes. It is one of the most remarkable discoveries which science owes to Euler. In 1761 he gave to the world the complete integration of an equation of two terms, each an elliptic function of the first or second order, not separately integrable.

This integration introduced an arbitrary constant in the form of a third function, related to the first two by a given equation between the amplitudes of the three.

In 1775 Landen, an English mathematician, published his celebrated theorem showing that any arc of a hyperbola may be measured by two arcs of an ellipse, an important element of the theory of Elliptic Functions, but *then* an isolated result. The great problem of comparison of Elliptic Functions of different moduli remained unsolved, though Euler, in a measure, exhausted the comparison of functions of the same modulus. It was completed in 1784 by Lagrange, and for the computation

* Condensed from an article by Rev. Henry Moseley, M.A., F.R.S., Prof. of Nat. Phil. and Ast., King's College, London.

of numerical results leaves little to be desired. The value of a function may be determined by it, in terms of increasing or diminishing moduli, until at length it depends upon a function having a modulus of zero, or unity.

For all practical purposes this was sufficient. The enormous task of calculating tables was undertaken by Legendre. His labors did not end here, however. There is none of the discoveries of his predecessors which has not received some perfection at his hands; and it was he who first supplied to the whole that connection and arrangement which have made it an independent science.

The theory of Elliptic Integrals remained at a standstill from 1786, the year when Legendre took it up, until the year 1827, when the second volume of his *Traité des Fonctions Elliptiques* appeared. Scarcely so, however, when there appeared the researches of Jacobi, a Professor of Mathematics in Königsberg, in the 123d number of the *Journal of Schumacher*, and those of Abel, Professor of Mathematics at Christiania, in the 3d number of *Crelle's Journal* for 1827.

These publications put the theory of Elliptic Functions upon an entirely new basis. The researches of Jacobi have for their principal object the development of that general relation of functions of the first order having different moduli, of which the scales of Lefschetz and Legendre are particular cases.

It was to Abel that the idea first occurred of treating the Elliptic Integral as a function of its amplitude. Proceeding from this new point of view, he embraced in his speculations all the principal results of Jacobi. Having undertaken to develop the principle upon which rests the fundamental proposition of Euler establishing an algebraic relation between three functions which have the same moduli, dependent upon a certain relation of their amplitudes, he has extended it from three to an indefinite number of functions; and from Elliptic Functions to an infinite number of other functions embraced under an indefinite number of classes, of which that of Elliptic Func-

tions is but one ; and each class having a division analogous to that of Elliptic Functions into three orders having common properties.

The discovery of Abel is of infinite moment as presenting the first step of approach towards a more complete theory of the infinite class of ultra elliptic functions, destined probably ere long to constitute one of the most important of the branches of transcendental analysis, and to include among the integrals of which it effects the solution some of those which at present arrest the researches of the philosopher in the very elements of physics.

CHAPTER I.

ELLIPTIC INTEGRALS.

THE integration of irrational expressions of the form

$$Xdx \sqrt{A + Bx + Cx^2},$$

or

$$\frac{Xdx}{\sqrt{A + Bx + Cx^2}},$$

X being a rational function of x , is fully illustrated in most elementary works on Integral Calculus, and shown to depend upon the transcendentials known as logarithms and circular functions, which can be calculated by the proper logarithmic and trigonometric tables.

When, however, we undertake to integrate irrational expressions containing higher powers of x than the square, we meet with insurmountable difficulties. This arises from the fact that the integral sought depends upon a new set of transcendentials, to which has been given the name of *elliptic functions*, and whose characteristics we will learn hereafter.

The name of Elliptic Integrals has been given to the simple integral forms to which can be reduced all integrals of the form

$$(I) \quad V = \int F(X, R)dx,$$

where $F(X, R)$ designates a rational function of x and R , and R represents a radical of the form

$$R = \sqrt{Ax^4 + Bx^3 + Cx^2 + Dx + E},$$

where A, B, C, D, E indicate constant coefficients.

We will show presently that all cases of Eq. (1) can be reduced to the three typical forms

$$(2) \quad \begin{aligned} & \int_0^x \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}}, \\ & \int_0^x \frac{x^2 dx}{\sqrt{(1-x^2)(1-k^2x^2)}}, \\ & \int_0^x \frac{dx}{(x^2+a)\sqrt{(1-x^2)(1-k^2x^2)}}, \end{aligned}$$

which are called elliptic integrals of the first, second, and third order.

Why they are called *Elliptic* Integrals we will learn further on. The transcendental functions which depend upon these integrals, and which will be discussed in Chapter IV, are called *Elliptic Functions*.

The most general form of Eq. (1) is

$$(3) \quad V = \int \frac{A + BR}{C + DR} dx;$$

where A, B, C , and D stand for rational integral functions of x .

$\frac{A + BR}{C + DR}$ can be written

$$\begin{aligned} \frac{A + BR}{C + DR} &= \frac{AC - BDR^2}{C^2 - D^2R^2} - \frac{(AD - CB)R^2}{C^2 - D^2R^2} \cdot \frac{1}{R} \\ &= N - \frac{P}{R}; \end{aligned}$$

N and P being rational integral functions of x . Whence Eq. (3) becomes

$$(4) \quad V = \int N dx - \int \frac{P dx}{R}.$$

Eq. (4) shows that the most general form of V can be made to depend upon the expressions

$$(5) \quad V' = \int \frac{P dx}{R},$$

and

$$\int N dx.$$

This last form is rational, and needs no discussion here. We can write

$$\begin{aligned} P &= \frac{G_0 + G_1 x + G_2 x^2 + \dots}{H_0 + H_1 x + H_2 x^2 + \dots} \\ &= \frac{G_0 + G_2 x^2 + G_4 x^4 + \dots + (G_1 + G_3 x^2 + \dots)x}{H_0 + H_2 x^2 + H_4 x^4 + \dots + (H_1 + H_3 x^2 + \dots)x}. \end{aligned}$$

Multiplying both numerator and denominator by

$$H_0 + H_2 x^2 + H_4 x^4 + \dots - (H_1 + H_3 x^2 + H_5 x^4 + \dots)x,$$

we have a new numerator which contains only powers of x^2 . The result takes the following form:

$$\begin{aligned} P &= \frac{M_0 + M_2 x^2 + M_4 x^4 + \dots + (M_1 + M_3 x^2 + M_5 x^4 + \dots)x}{N_0 + N_2 x^2 + N_4 x^4 + N_6 x^6 + \dots} \\ &= \Phi(x^2) + \Psi(x^2) \cdot x. \end{aligned}$$

Equation (5) thus becomes

$$(6) \quad V' = \int \frac{\Phi(x^2) dx}{R} + \int \frac{\Psi(x^2) \cdot x \cdot dx}{R}.$$

We shall see presently that R can always be assumed to be of the form

$$\sqrt{(1-x^2)(1-k^2x^2)}.$$

Therefore, putting $x^2 = z$, the second integral in Eq. (6) takes the form

$$\frac{1}{2} \int \frac{\Psi(z) \cdot dz}{\sqrt{(1-z)(1-k^2z)}},$$

which can be integrated by the well-known methods of Integral Calculus, resulting in logarithmic and circular transcendental.

There remains, therefore, only the form

$$\int \frac{\Phi(x^2)dx}{R}$$

to be determined.

We will now show that R can always be assumed to be in the form

$$\sqrt{(1-x^2)(1-k^2x^2)}.$$

We have

$$\begin{aligned} R &= \sqrt{Ax^4 + Bx^3 + Cx^2 + Dx + E} \\ &= \sqrt{G(x-a)(x-b)(x-c)(x-d)}, \end{aligned}$$

a, b, c , and d being the roots of the polynomial of the fourth degree, and G any number, real or imaginary, depending upon the coefficients in the given polynomial.

Substituting in equation (1).

$$x = \frac{p + qy}{1 + y},$$

we have

$$(7) \quad V = \int \phi(y, \rho) dy,$$

ρ designating the radical

$$\rho = \sqrt{G[p-a+(q-a)y][p-b+(q-b)y][p-c+(q-c)y] \dots}$$

In order that the odd powers of y under the radical may disappear we must have their coefficients equal to zero; i.e.,

$$\begin{aligned}(p-a)(q-b) + (p-b)(q-a) &= 0, \\ (p-c)(q-d) + (p-d)(q-c) &= 0;\end{aligned}$$

whence

$$\begin{aligned}2pq - (p+q)(a+b) + 2ab &= 0, \\ 2pq - (p+q)(c+d) + 2cd &= 0,\end{aligned}$$

and

$$(8) \quad \begin{cases} pq = \frac{ab(c+d) - cd(a+b)}{a+b-(c+d)}, \\ p+q = \frac{2ab-2cd}{a+b-(c+d)}. \end{cases}$$

Equation (8) shows that p and q are real quantities, whether the roots a, b, c , and d are real or imaginary; a, b , and c, d being the conjugate pairs.

Hence equation (1) can always be reduced to the form of equation (7), which contains only the second and fourth powers of the variable.

This transformation seems to fail when $a+b-(c+d)=0$; but in that case we have

$$R = \sqrt{G[x^2 - (a+b)x + ab][x^2 - (a+b)x + cd]},$$

and substituting

$$x = y - \frac{a+b}{2}$$

will cause the odd powers of y to disappear as before.

If the radical should have the form

$$\sqrt{G(x-a)(x-b)(x-c)},$$

placing $x = y^2 + a$, we get

$$V = \int \phi(y, \rho) dy,$$

$$\rho = \sqrt{G(y^2 + a - b)(y^2 + a - c)},$$

ϕ designating a rational function of y and ρ .

Thus all integrals of the form contained in equation (1), in which R stands for a quadratic surd of the third or fourth degree, can be reduced to the form

$$(9) \quad V = \int \phi(x, R) dx,$$

R designating a radical of the form

$$\sqrt{G(1 + mx^2)(1 + nx^2)},$$

m and n designating constants.

It is evident that if we put

$$x' = x \sqrt{-m}, \quad k^2 = -\frac{n}{m},$$

we can reduce the radical to the form

$$\sqrt{(1 - x'^2)(1 - k^2 x'^2)}.$$

We shall see later on that the quantity k^2 , to which has been given the name *modulus*, can always be considered real and less than unity.

Combining these results with equation (6), we see that the integration of equation (1) depends finally upon the integration of the expression

$$(10) \quad V'' = \int \frac{\phi(x^2) dx}{\sqrt{(1 - x^2)(1 - k^2 x^2)}} = \int \frac{\phi(x^2) dx}{R}.$$

The most general form of $\phi(x^2)$ is

$$\begin{aligned}\phi(x^2) &= \frac{M_0 + M_2x^2 + M_4x^4 + \dots}{N_0 + N_2x^2 + N_4x^4 + \dots} \\ &= P_0 + P_2x^2 + P_4x^4 + P_6x^6 + \dots \\ &\quad + \Sigma \frac{L}{(x^2 + a)^n}.\end{aligned}$$

Hence

$$(11) \quad V'' = \Sigma P \int \frac{x^{2m} dx}{R} + \Sigma L \int \frac{dx}{(x^2 + a)^n R}.$$

But $\int \frac{x^{2m} dx}{R}$ depends upon $\int \frac{dx}{R}$ and $\int \frac{x^2 dx}{R}$, which can be shown as follows:

Differentiating Rx^{2m-3} , we have

$$\begin{aligned}d[x^{2m-3}R] &= d[x^{2m-3} \sqrt{\alpha + \beta x^2 + \gamma x^4}] \\ &= (2m-3)x^{2m-4}dx \sqrt{\alpha + \beta x^2 + \gamma x^4} \\ &\quad + \frac{x^{2m-3}(\beta x + 2\gamma x^3)dx}{\sqrt{\alpha + \beta x^2 + \gamma x^4}}.\end{aligned}$$

Integrating and collecting, we get

$$\begin{aligned}Rx^{2m-3} &= (2m-3)\alpha \int \frac{x^{2m-4}dx}{R} + (2m-2)\beta \int \frac{x^{2m-2}dx}{R} \\ &\quad + (2m-1)\gamma \int \frac{x^{2m}dx}{R} \\ (12) \quad &= \alpha' \int \frac{x^{2m-4}dx}{R} + \beta' \int \frac{x^{2m-2}dx}{R} + \gamma' \int \frac{x^{2m}dx}{R}.\end{aligned}$$

Whence we get, by taking $m = 2$,

$$(13) \quad Rx = \alpha \int \frac{dx}{R} + \beta \int \frac{x^2 dx}{R} + \gamma \int \frac{x^4 dx}{R},$$

which shows that the general expression $\int \frac{x^{2m} dx}{R}$ can be found by successive calculations, when we are able to integrate the expressions

$$\int \frac{dx}{R} \quad \text{and} \quad \int \frac{x^2 dx}{R},$$

the first and second of equation (2).

We will now consider the second class of terms in eq. (11), viz., $\frac{Ldx}{(x^2 + a)^n R}$.

This second term is as follows :

$$(14) \quad \Sigma \int \frac{L}{(x^2 + a)^n R} = \int \frac{A dx}{(x^2 + a)^n R} + \int \frac{B dx}{(x^2 + a)^{n-1} R} \\ + \int \frac{C dx}{(x^2 + a)^{n-2} R} + \dots$$

Each of these terms can be shown to depend ultimately upon terms of the form

$$\frac{x^2 dx}{R}, \quad \frac{dx}{R}, \quad \text{and} \quad \frac{dx}{(x^2 + a)R}.$$

The two former will be recognized as the two ultimate forms already discussed, the first and second of equation (2). The third is the third one of equation (2).

This dependence of equation (14) can be shown as follows :

We have

$$\begin{aligned} d\left[\frac{xR}{(x^2+a)^{n-1}}\right] &= \frac{(x^2+a)^{n-1}(xdR+Rdx)-2x^2R(n+1)(x^2+a)^{n-2}dx}{(x^2+a)^{2n-2}} \\ &= \frac{(x^2+a)(xdR+Rdx)-2x^2R(n-1)dx}{(x^2+a)^n}. \end{aligned}$$

Substituting the value of

$$R = \sqrt{\alpha + \beta x^2 + \gamma x^4} \quad \text{and} \quad dR = (\beta x + 2\gamma x^3) \frac{dx}{R},$$

we get

$$\begin{aligned} d\left[\frac{xR}{(x^2+a)^{n-1}}\right] &= \\ \frac{(x^2+a)(\beta x^2+2\gamma x^4+\alpha+\beta x^2+\gamma x^4)-2x^2(n-1)(\alpha+\beta x^2+\gamma x^4)}{(x^2+a)^n} \cdot \frac{dx}{R} \\ &= \frac{-2(n-1)\gamma \left| \begin{array}{c} 3\gamma \mid x^6 \\ +2\beta \mid x^4 \\ +3a\gamma \mid \\ +2a\beta \mid x^2+a\alpha \end{array} \right.}{(x^2+a)^n} \cdot \frac{dx}{R} \\ &= \frac{-(2n-5)\gamma x^6-(2n-4)\beta \left| \begin{array}{c} x^4-(2n-3)\alpha \mid x^2+a\alpha \\ +3a\gamma \mid \\ -2a\beta \mid \end{array} \right.}{(x^2+a)^n} \cdot \frac{dx}{R}; \end{aligned}$$

or, by substituting in the numerator $x^2 = z - a$,

$$\begin{aligned} &-(2n-5)\gamma z^3+(2n-5)3a\gamma \left| \begin{array}{c} z^2-(2n-5)3a^2\gamma \mid z+(2n-5)a^3\gamma \\ -(2n-4)\beta \mid + (2n-4)2a\beta \mid -(2n-4)a^2\beta \\ +3a\gamma \mid -6a^2\gamma \mid +3a^3\gamma \\ - (2n-3)\alpha \mid + (2n-3)a\alpha \\ +2a\beta \mid -2a^2\beta \\ +a\alpha \end{array} \right. \\ &= \frac{\quad}{(x^2+a)^n} \cdot \frac{dx}{R}; \end{aligned}$$

or, after resubstituting $z = x^2 + a$, and integrating,

$$\begin{aligned}
 (15) \quad \frac{xR}{(x^2 + a)^{n-1}} &= -(2n-5)\gamma \int \frac{dx}{(x^2 + a)^{n-3}R} \\
 &\quad - (2n-4)(\beta - 3a\gamma) \int \frac{dx}{(x^2 + a)^{n-2}R} \\
 &\quad - (2n-3)(3a^2\gamma - 2a\beta + \alpha) \int \frac{dx}{(x^2 + a)^{n-1}R} \\
 &\quad + (2n-2)(a^2\gamma - a^2\beta + a\alpha) \int \frac{dx}{(x^2 + a)^n R} \\
 &= \alpha_1 \int \frac{dx}{(x^2 + a)^{n-3}R} + \beta_1 \int \frac{dx}{(x^2 + a)^{n-2}R} + \gamma_1 \int \frac{dx}{(x^2 + a)^{n-1}R} \\
 &\quad + \delta_1 \int \frac{dx}{(x^2 + a)^n R}.
 \end{aligned}$$

Making $n = 2$, we have

$$\begin{aligned}
 (16) \quad \frac{xR}{(x^2 + a)^{-1}} &= \alpha_1 \int \frac{(x^2 + a)dx}{R} + \beta_1 \int \frac{dx}{R} + \gamma_1 \int \frac{dx}{(x^2 + a)R} \\
 &\quad + \delta_1 \int \frac{dx}{(x^2 + a)^2 R}.
 \end{aligned}$$

Equation (16) shows that

$$\int \frac{dx}{(x^2 + a)^2 R}$$

depends upon the three forms

$$\int \frac{x^2 dx}{R}, \quad \int \frac{dx}{R}, \quad \text{and} \quad \int \frac{dx}{(x^2 + a)R},$$

the three types of equation (2), and equation (15) shows that the general form

$$\int \frac{dx}{(x^2 + a)^n R}$$

depends ultimately upon the same three types.

We have now discussed every form which the general equation (1) can assume, and shown that they all depend ultimately upon one or more of the three types contained in equation (2).

These three types are called the three Elliptic Integrals of the first, second, and third kind, respectively.

Legendre puts $x = \sin \phi$, and reduces the three integrals to the following forms:

$$(17) \quad F(k, \phi) = \int_0^\phi \frac{d\phi}{\sqrt{1 - k^2 \sin^2 \phi}};$$

$$\frac{1}{k^2} \int_0^\phi \frac{d\phi}{\sqrt{1 - k^2 \sin^2 \phi}} - \frac{1}{k^2} \int_0^\phi \sqrt{1 - k^2 \sin^2 \phi} \cdot d\phi;$$

$$(18) \quad \Pi(n, k, \phi) = \int_0^\phi \frac{d\phi}{(1 - n \sin^2 \phi) \sqrt{1 - k^2 \sin^2 \phi}};$$

the first being Legendre's integral of the first kind; the form

$$(19) \quad E(k, \phi) = \int_0^\phi \sqrt{1 - k^2 \sin^2 \phi} \cdot d\phi$$

being the integral of the second kind; and the third one being the integral of the third kind.

The form of the integral of the second kind shows why they are called Elliptic Integrals, the arc of an elliptic quadrant being equal to

$$a \int_0^{\frac{\pi}{2}} \sqrt{1 - e^2 \sin^2 \phi} \cdot d\phi,$$

ϕ being the complement of the eccentric angle.

By easy substitutions, we get from Eqs. (17), (18), and (19) the following solutions:

$$\int_0^\phi \frac{\sin^2 \phi}{\Delta} d\phi = \frac{F - E}{k^2};$$

$$\int_0^\phi \frac{\cos^2 \phi}{\Delta} d\phi = \frac{E - (1 - k^2)F}{k^2};$$

$$\int_0^\phi \frac{\tan^2 \phi}{\Delta} d\phi = \frac{\Delta \tan \phi - E}{1 - k^2};$$

$$\int_0^\phi \frac{\sec^2 \phi}{\Delta} d\phi = \frac{\Delta \tan \phi + (1 - k^2)F - E}{1 - k^2};$$

$$\int_0^\phi \frac{1}{\Delta^3} d\phi = \frac{1}{1 - k^2} \left(E - \frac{k^2 \sin \phi \cos \phi}{\Delta} \right);$$

$$\int_0^\phi \frac{\sin^2 \phi}{\Delta^3} d\phi = \frac{1}{1 - k^2} \left(\frac{E - (1 - k^2)F}{k^2} - \frac{\sin \phi \cos \phi}{\Delta} \right);$$

$$\int_0^\phi \frac{\cos^2 \phi}{\Delta^3} d\phi = \frac{F - E}{k^2} + \frac{\sin \phi \cos \phi}{\Delta}.$$

CHAPTER II.

ELLIPTIC FUNCTIONS.

LET
$$u = \int_0^\phi \frac{d\phi}{\sqrt{1 - k^2 \sin^2 \phi}}.$$

ϕ^* is called the *amplitude* corresponding to the *argument* u , and is written

$$\phi = \text{am}(u, k) = \text{am } u.$$

The quantity k is called the *modulus*, and the expression $\sqrt{1 - k^2 \sin^2 \phi}$ is written *

$$\sqrt{1 - k^2 \sin^2 \phi} = \Delta \text{ am } u = \Delta \phi,$$

and is called the *delta function* of the amplitude o. u , or *delta of ϕ* , or simply *delta ϕ* .

u can be written

$$u = F(k, \phi).$$

The following abbreviations are used:

$$\sin \phi = \sin \text{ am } u = \text{sn } \dagger u;$$

$$\cos \phi = \cos \text{ am } u = \text{cn } \dagger u;$$

$$\Delta \phi = \Delta \text{ am } u = \text{dn } \dagger u = \Delta u;$$

$$\tan \phi = \tan \text{ am } u = \text{tn } u.$$

Let ϕ and ψ be any two arbitrary angles, and put

$$\phi = \text{am } u;$$


$$\psi = \text{am } v.$$

* Legendre.

† Gudermann, in his "Theorie der Modularfunctionen": Crelle's Journal, Bd. 18.

In the spherical triangle ABC we have from Trigonometry, c and C being constant,

$$\frac{d\phi}{\cos B} + \frac{d\psi}{\cos A} = 0.$$



Since C and c are constant, denoting by k an arbitrary constant, we have

$$(1) \quad \frac{\sin C}{\sin \mu} = k.$$

But

$$\sin A = \sin \psi \frac{\sin B}{\sin \phi} = \sin \psi \frac{\sin C}{\sin \mu} = k \sin \psi.$$

Whence

$$\cos A = \sqrt{1 - \sin^2 A} = \sqrt{1 - k^2 \sin^2 \psi}.$$

In the same manner

$$\cos B = \sqrt{1 - \sin^2 B} = \sqrt{1 - k^2 \sin^2 \phi}.$$

Substituting these values, we get

$$(2) \quad \frac{d\phi}{\sqrt{1 - k^2 \sin^2 \phi}} + \frac{d\psi}{\sqrt{1 - k^2 \sin^2 \psi}} = 0.$$

Integrating this, there results

$$(3) \quad \int_0^\phi \frac{d\phi}{\sqrt{1 - k^2 \sin^2 \phi}} + \int_0^\psi \frac{d\psi}{\sqrt{1 - k^2 \sin^2 \psi}} = \text{const.}$$

When $\phi = 0$, we have $\psi = \mu$, and therefore the constant must be of the form

$$\int_0^\mu \frac{d\phi}{\sqrt{1 - k^2 \sin^2 \phi}},$$

whence

$$(4) \int_0^\phi \frac{d\phi}{\sqrt{1 - k^2 \sin^2 \phi}} + \int_0^\psi \frac{d\psi}{\sqrt{1 - k^2 \sin^2 \psi}} = \int_0^\mu \frac{d\phi}{\sqrt{1 - k^2 \sin^2 \phi}},$$

or

$$u + v = m;$$

and evidently the amplitudes ϕ , ψ , and μ can be considered as the three sides of a spherical triangle, and the relations between the sides of this spherical triangle will be the same as those between ϕ , ψ , and μ .


But the sides of this triangle have imposed upon them the condition

$$\frac{\sin C}{\sin \mu} = k;$$

and since $k < 1$, we must have $\mu > C$, which requires that one of the angles of the triangle shall be obtuse and the other two acute.

In the figure, let C be an acute angle of the triangle ABC , and PQ the equatorial great circle of which C is the pole.

The arc PQ will be the measure of the angle C .



Let AG and AH be the arcs of two great circles perpendicular respectively to CQ and CP . They will of course be shorter than PQ . Hence $AB = \mu$ must intersect CQ in points between CG and HQ , since $\mu > (C = PQ)$. In any case either A or B will be obtuse according as B falls between QH or CG respectively; and

the other angle will be acute.

In the case where C is an obtuse angle, it will be easily seen that the angle at A must be acute, since the great circle AD , perpendicular to CP , intersects PQ in D , PD being a quadrant. The same remarks apply to the angle B . Hence, in either

case, one of the angles of the triangle is obtuse and the other two are acute, as a result of the condition

$$\frac{\sin C}{\sin \mu} = k < 1.$$

From Trigonometry we have

$$\cos \mu = \cos \phi \cos \psi + \sin \phi \sin \psi \cos C;$$

and since the angle C is obtuse,

$$\cos C = -\sqrt{1 - \sin^2 C} = -\sqrt{1 - k^2 \sin^2 \mu},$$

and

$$(5) \quad \cos \mu = \cos \phi \cos \psi - \sin \phi \sin \psi \sqrt{1 - k^2 \sin^2 \mu},$$

the relation sought.

The spherical triangle likewise gives the following relations between the sides:

$$(5)^* \quad \begin{cases} \cos \phi = \cos \mu \cos \psi + \sin \mu \sin \psi \sqrt{1 - k^2 \sin^2 \phi}; \\ \cos \psi = \cos \mu \cos \phi + \sin \mu \sin \phi \sqrt{1 - k^2 \sin^2 \psi}. \end{cases}$$

These give, by eliminating $\cos \mu$,

$$\sin \mu = \frac{\cos^2 \psi - \cos^2 \phi}{\sin \phi \cos \psi \Delta \psi - \sin \psi \cos \phi \Delta \phi};$$

which, after multiplying by the sum of the terms in the denominator and substituting $\cos^2 = 1 - \sin^2$, can be written


$$\sin \mu = \frac{(\sin^2 \phi - \sin^2 \psi)(\sin \phi \cos \psi \Delta \psi + \sin \psi \cos \phi \Delta \phi)}{\sin^2 \phi \cos^2 \psi \Delta^2 \psi - \sin^2 \psi \cos^2 \phi \Delta^2 \phi}.$$

Since the denominator can be written

$$(6) \quad \sin \mu = \frac{(\sin^2 \phi - \sin^2 \psi)(1 - k^2 \sin^2 \phi \sin^2 \psi),}{1 - k^2 \sin^2 \phi \sin^2 \psi}.$$

In a similar manner we get

$$(6)^* \quad \begin{cases} \cos \mu = \frac{\cos \phi \cos \psi - \sin \phi \sin \psi \Delta \phi \Delta \psi}{1 - k^2 \sin^2 \phi \sin^2 \psi}; \\ \Delta \mu = \frac{\Delta \phi \Delta \psi - k^2 \sin \phi \sin \psi \cos \phi \cos \psi}{1 - k^2 \sin^2 \phi \sin^2 \psi}. \end{cases}$$



These equations can also be written as follows:

$$(7) \begin{cases} \sin \operatorname{am} (u \pm v) = \frac{\sin \operatorname{am} u \cos \operatorname{am} v \Delta \operatorname{am} v \pm \sin \operatorname{am} v \cos \operatorname{am} u \Delta \operatorname{am} u}{1 - k^2 \sin^2 \operatorname{am} u \sin^2 \operatorname{am} v}; \\ \cos \operatorname{am} (u \pm v) = \frac{\cos \operatorname{am} u \cos \operatorname{am} v \mp \sin \operatorname{am} u \sin \operatorname{am} v \Delta \operatorname{am} u \Delta \operatorname{am} v}{1 - k^2 \sin^2 \operatorname{am} u \sin^2 \operatorname{am} v}; \\ \Delta \operatorname{am} (u \pm v) = \frac{\Delta \operatorname{am} u \Delta \operatorname{am} v \mp k^2 \sin \operatorname{am} u \sin \operatorname{am} v \cos \operatorname{am} u \cos \operatorname{am} v}{1 - k^2 \sin^2 \operatorname{am} u \sin^2 \operatorname{am} v}; \end{cases}$$

or

$$(8) \begin{cases} \operatorname{sn} (u \pm v) = \frac{\operatorname{sn} u \operatorname{cn} v \operatorname{dn} v \pm \operatorname{sn} v \operatorname{cn} u \operatorname{dn} u}{1 - k^2 \operatorname{sn}^2 u \operatorname{sn}^2 v}; \\ \operatorname{cn} (u \pm v) = \frac{\operatorname{cn} u \operatorname{cn} v \mp \operatorname{sn} u \operatorname{sn} v \operatorname{dn} u \operatorname{dn} v}{1 - k^2 \operatorname{sn}^2 u \operatorname{sn}^2 v}; \\ \operatorname{dn} (u \pm v) = \frac{\operatorname{dn} u \operatorname{dn} v \mp k^2 \operatorname{sn} u \operatorname{sn} v \operatorname{cn} u \operatorname{cn} v}{1 - k^2 \operatorname{sn}^2 u \operatorname{sn}^2 v}. \end{cases}$$

Making $u = v$, we get from the upper sign

$$(9) \begin{cases} \operatorname{sn} 2u = \frac{2 \operatorname{sn} u \operatorname{cn} u \operatorname{dn} u}{1 - k^2 \operatorname{sn}^4 u}; \\ \operatorname{cn} 2u = \frac{\operatorname{cn}^2 u - \operatorname{sn}^2 u \operatorname{dn}^2 u}{1 - k^2 \operatorname{sn}^4 u} = \frac{1 - 2 \operatorname{sn}^2 u + k^2 \operatorname{sn}^4 u}{1 - k^2 \operatorname{sn}^4 u}; \\ \operatorname{dn} 2u = \frac{\operatorname{dn}^2 u - k^2 \operatorname{sn}^2 u \operatorname{cn}^2 u}{1 - k^2 \operatorname{sn}^4 u} = \frac{1 - 2k^2 \operatorname{sn}^2 u + k^2 \operatorname{sn}^4 u}{1 - k^2 \operatorname{sn}^4 u}. \end{cases}$$

From these

$$(10) \begin{cases} 1 - \operatorname{cn} 2u = \frac{2 \operatorname{cn}^2 u \operatorname{dn}^2 u}{1 - k^2 \operatorname{sn}^4 u}; \\ 1 + \operatorname{cn} 2u = \frac{2 \operatorname{cn}^2 u}{1 - k^2 \operatorname{sn}^4 u}; \\ 1 - \operatorname{dn} u = \frac{2k^2 \operatorname{sn}^2 u \operatorname{cn}^2 u}{1 - k^2 \operatorname{sn}^4 u}; \\ 1 + \operatorname{dn} u = \frac{2 \operatorname{dn}^2 u}{1 - k^2 \operatorname{sn}^4 u}; \end{cases}$$

and therefore

$$(11) \quad \begin{cases} \operatorname{sn}^2 u = \frac{1 - \operatorname{cn} 2u}{1 + \operatorname{dn} 2u}; \\ \operatorname{cn}^2 u = \frac{\operatorname{dn} 2u + \operatorname{cn} 2u}{1 + \operatorname{dn} 2u}; \\ \operatorname{dn}^2 u = \frac{1 - k^2 + \operatorname{dn} 2u + k^2 \operatorname{cn} 2u}{1 + \operatorname{dn} 2u}; \end{cases}$$

and by analogy

$$(12) \quad \begin{cases} \operatorname{sn} \frac{u}{2} = \sqrt{\frac{1 - \operatorname{cn} u}{1 + \operatorname{dn} u}}; \\ \operatorname{cn} \frac{u}{2} = \sqrt{\frac{\operatorname{cn} u + \operatorname{dn} u}{1 + \operatorname{dn} u}}; \\ \operatorname{dn} \frac{u}{2} = \sqrt{\frac{1 - k^2 + \operatorname{dn} u + k^2 \operatorname{cn} u}{1 + \operatorname{dn} u}}. \end{cases}$$

In equations (7) making $u = v$, and taking the lower sign, we have

$$(13) \quad \begin{cases} \operatorname{sn} 0 = 0; \\ \operatorname{cn} 0 = 1; \\ \operatorname{dn} 0 = 1. \end{cases}$$

Likewise, we get by making $u = 0$,

$$(14) \quad \begin{cases} \operatorname{sn} (-u) = -\operatorname{sn} u; \\ \operatorname{cn} (-u) = +\operatorname{cn} u; \\ \operatorname{dn} (-u) = \operatorname{dn} u. \end{cases}$$

CHAPTER III.

PERIODICITY OF THE FUNCTIONS.

WHEN the elliptic integral

$$\int_0^\phi \frac{d\phi}{\sqrt{1 - k^2 \sin^2 \phi}}$$

has for its amplitude $\frac{\pi}{2}$, it is called, following the notation of Legendre, the *complete* function, and is indicated by K , thus :

$$K = \int_0^{\frac{\pi}{2}} \frac{d\phi}{\sqrt{1 - k^2 \sin^2 \phi}}.$$

When k becomes the complementary modulus, k' , (see eq. 4, Chap. IV,) the corresponding complete function is indicated by K' , thus :

$$K' = \int_0^{\frac{\pi}{2}} \frac{d\phi}{\sqrt{1 - k'^2 \sin^2 \phi}}.$$

From these, evidently,

$$\text{am } (K, k) = \frac{\pi}{2}, \quad \text{am } (K', k') = \frac{\pi}{2}.$$

$$(I) \quad \begin{cases} \text{sn } (K, k) = 1; \\ \text{cn } (K, k) = 0; \\ \text{dn } (K, k) = k'. \end{cases}$$

From eqs. (7), (8), and (9), Chap. II, we have, by the substitution of the values of $\text{sn}(K) = 1$, $\text{cn}(K) = 0$, $\text{dn}(K) = k'$,

$$(2) \quad \begin{cases} \text{sn } 2K = 0; \\ \text{cn } 2K = -1; \\ \text{dn } 2K = 1. \end{cases}$$

These equations, by means of (1), (2), and (3) of Chap. II, give

$$(3) \quad \begin{cases} \text{sn}(u + 2K) = -\text{sn } u; \\ \text{cn}(u + 2K) = -\text{cn } u; \\ \text{dn}(u + 2K) = \text{dn } u; \end{cases}$$

and these, by changing u into $u + 2K$, give

$$(4) \quad \begin{cases} \text{sn}(u + 4K) = \text{sn } u; \\ \text{cn}(u + 4K) = \text{cn } u; \\ \text{dn}(u + 4K) = \text{dn } u. \end{cases}$$

From these equations it is seen that the elliptic functions sn , cn , dn , are periodic functions having for their period $4K$. Unlike the period of trigonometric functions, this period is not a fixed one, but depends upon the value of k , the modulus.

From the Integral Calculus we have

$$\begin{aligned} \int_0^{n\frac{\pi}{2}} \frac{d\phi}{\Delta\phi} &= \int_0^{\frac{\pi}{2}} \frac{d\phi}{\Delta\phi} + \int_{\frac{\pi}{2}}^{\pi} \frac{d\phi}{\Delta\phi} + \int_{\pi}^{\frac{3\pi}{2}} \frac{d\phi}{\Delta\phi} + \dots + \int_{(n-1)\frac{\pi}{2}}^{n\frac{\pi}{2}} \frac{d\phi}{\Delta\phi} \\ &= n \int_0^{\frac{\pi}{2}} \frac{d\phi}{\Delta\phi} = nK; \end{aligned}$$

from which we see that

$$n\frac{\pi}{2} = \text{am}(nK);$$

or, since $\frac{\pi}{2} = \text{am } K$,

$$\text{am } (nK) = n \cdot \text{am } K,$$

and

$$n\pi = \text{am } (2nK),$$

and also

$$n\pi = 2n \text{ am } K.$$

In the case of an Elliptic Integral with the arbitrary angle α , we can put

$$d = n\pi \pm \beta,$$

where β is an angle between 0 and $\frac{\pi}{2}$, the upper or the lower sign being taken according as $\frac{\pi}{2}$ is contained in α an even or an uneven number of times.

In the first case we have

$$\int_0^{n\pi+\beta} \frac{d\phi}{\Delta\phi} = \int_0^{n\pi} \frac{d\phi}{\Delta\phi} + \int_{n\pi}^{n\pi+\beta} \frac{d\phi}{\Delta\phi};$$

or, putting $\phi_1 = -n\pi$,

$$\int_0^{n\pi+\beta} \frac{d\phi}{\Delta\phi} = 2nK + \int_0^{\beta} \frac{d\phi_1}{\Delta\phi_1}.$$

In the second case

$$\int_0^{n\pi-\beta} \frac{d\phi}{\Delta\phi} = \int_0^{n\pi} \frac{d\phi}{\Delta\phi} - \int_{n\pi-\beta}^{n\pi} \frac{d\phi}{\Delta\phi};$$

or, putting $\phi_1 = n\pi - \phi$,

$$\int_0^{n\pi-\beta} \frac{d\phi}{\Delta\phi} = 2nK - \int_0^{\beta} \frac{d\phi_1}{\Delta\phi_1};$$

or in either case,

$$\int_0^{n\pi \pm \beta} \frac{d\phi}{\Delta\phi} = 2nK \pm \int_0^{\beta} \frac{d\phi_1}{\Delta\phi_1}.$$

Thus we see that the Integral with the general amplitude α can be made to depend upon the complete integral K and an Integral whose amplitude lies between 0 and $\frac{\pi}{2}$.

Put now

$$\int_0^{\beta} \frac{d\phi_1}{\Delta\phi_1} = u, \quad \beta = \text{am } u.$$

This gives

$$\int_0^{n\pi \pm \beta} \frac{d\phi}{\Delta\phi} = 2nK \pm u,$$

or

$$\text{am } (2nK \pm u) = n\pi \pm \beta$$

(5)

$$= n\pi \pm \text{am } u$$

(6)

$$= 2n \cdot \text{am } K \pm \text{am } u;$$

or, since

$$\text{am } (-z) = -\text{am } z,$$

$$\text{am } (u \pm 2nK) = \text{am } u \pm n\pi$$

$$= \text{am } u \pm 2n \cdot \text{am } K.$$

Taking the sine and cosine of both sides, we have

$$\text{sn } (u \pm 2nK) = \pm \text{sn } u;$$

$$\text{cn } (u \pm 2nK) = \pm \text{cn } u;$$

the upper or the lower sign being taken according as n is even or odd. By giving the proper values to n we can get the same results as in equations (3) and (4).

Putting $n = 1$ in eq. (5), we have

$$\text{sn } (2K - u) = \sin \pi \text{cn } u - \cos \pi \text{sn } u$$

(7)

$$= \text{sn } u.$$

Elliptic functions also have an imaginary period. In order to show this we will, in the integral

$$\int_0^\phi \frac{d\phi}{\Delta\phi},$$

assume the amplitude as imaginary. Put

$$\sin \phi = i \tan \psi.$$

From this we get

$$(8) \quad \begin{cases} \cos \phi = \frac{1}{\cos \psi}; \\ \Delta\phi = \frac{\sqrt{1 - k'^2 \sin^2 \psi}}{\cos \psi} = \frac{\Delta(\psi, k')}{\cos \psi}; \\ d\phi = i \frac{d\psi}{\cos \psi}. \end{cases}$$

From these, since ϕ and ψ vanish simultaneously, we easily get

$$\int_0^\phi \frac{d\phi}{\Delta\phi} = i \int_0^\psi \frac{d\psi}{\Delta(\psi, k')}.$$

Put

$$\int_0^\psi \frac{d\psi}{\Delta(\psi, k')} = u \quad \text{and} \quad \psi = \text{am}(u, k'),$$

whence

$$\int_0^\phi \frac{d\phi}{\Delta\phi} = iu \quad \text{and} \quad \phi = \text{am}(iu);$$

and these substituted in Eq. (8) give

$$(9) \quad \begin{cases} \text{sn } iu = i \text{tn}(u, k'); \\ \text{cn } iu = \frac{1}{\text{cn}(u, k')}; \\ \text{dn } iu = \frac{\text{dn}(u, k')}{\text{cn}(u, k')}. \end{cases}$$

By assuming

$$\int_0^\psi \frac{d\psi}{\Delta(\psi, k')} = iu \quad \text{and} \quad \int_0^\phi \frac{d\phi}{\Delta\phi} = -u,$$

we get

$$\operatorname{sn}(-u) = i \operatorname{tn}(iu, k'),$$

$$\operatorname{cn}(-u) = \frac{1}{\operatorname{cn}(iu, k')},$$

$$\operatorname{dn}(-u) = \frac{\operatorname{dn}(iu, k')}{\operatorname{cn}(iu, k')};$$

or, from eq. (14), Chap. II,

$$(10) \quad \begin{cases} \operatorname{sn} u = -i \operatorname{tn}(iu, k'); \\ \operatorname{cn} u = \frac{1}{\operatorname{cn}(iu, k')}; \\ \operatorname{dn} u = \frac{\operatorname{dn}(iu, k')}{\operatorname{cn}(iu, k')}. \end{cases}$$

From eqs. (7), Chap. II, making $v = K$, we get, since $\operatorname{sn} K = 1$, $\operatorname{cn} K = 0$, $\operatorname{dn} K = k'$,

$$(11) \quad \begin{cases} \operatorname{sn}(u \pm K) = \pm \frac{\operatorname{cn} u \operatorname{dn} u}{1 - k'^2 \operatorname{sn}^2 u} = \pm \frac{\operatorname{cn} u}{\operatorname{dn} u}; \\ \operatorname{cn}(u \pm K) = \frac{\mp \operatorname{sn} u \operatorname{dn} u k'}{\operatorname{dn}^2 u} = \mp \frac{k' \operatorname{sn} u}{\operatorname{dn} u}; \\ \operatorname{dn}(u \pm K) = \pm \frac{k'}{\operatorname{dn} u}. \end{cases}$$

In these equations, changing u into iu , we get, by means of eqs. (9),

$$(12) \quad \begin{cases} \operatorname{sn}(iu \pm K) = \pm \frac{1}{\operatorname{dn}(u, k')}; \\ \operatorname{cn}(iu \pm K) = \mp \frac{ik' \operatorname{sn}(u, k')}{\operatorname{dn}(u, k')}; \\ \operatorname{dn}(iu \pm K) = \pm \frac{k' \operatorname{cn}(u, k')}{\operatorname{dn}(u, k')}. \end{cases}$$

Putting now in eqs. (9) $u \pm K'$ instead of u , and making use of eqs. (10), and interchanging k and k' , we have

$$(13) \quad \begin{cases} \operatorname{sn}(iu \pm iK') = -\frac{i \operatorname{cn}(u, k')}{k \operatorname{sn}(u, k')}; \\ \operatorname{cn}(iu \pm iK') = \mp \frac{\operatorname{dn}(u, k')}{k \operatorname{sn}(u, k')}; \\ \operatorname{dn}(iu \pm iK') = \mp \frac{1}{\operatorname{sn}(u, k')}. \end{cases}$$

Substituting in these $-iu$ in place of u , we get, by means of eqs. (9) and eqs. (14) of Chap. II,

$$(14) \quad \begin{cases} \operatorname{sn}(u \pm iK') = \frac{1}{k \operatorname{sn} u}; \\ \operatorname{cn}(u \pm iK') = \mp \frac{i \operatorname{dn} u}{k \operatorname{sn} u}; \\ \operatorname{dn}(u \pm iK') = \mp i \cot \operatorname{am} u. \end{cases}$$

In these equations, putting $u + K$ in place of u , we get

$$(15) \quad \begin{cases} \operatorname{sn}(u + K \pm iK') = \pm \frac{\operatorname{dn} u}{k \operatorname{cn} u}; \\ \operatorname{cn}(u + K \pm iK') = \mp \frac{ik'}{k \operatorname{cn} u}; \\ \operatorname{dn}(u + K \pm iK') = \pm ik' \operatorname{tn} u. \end{cases}$$

Whence for $u = 0$ we get

$$(16) \quad \begin{cases} \operatorname{sn}(K \pm iK') = \frac{1}{k}; \\ \operatorname{cn}(K \pm iK') = \mp \frac{ik'}{k}; \\ \operatorname{dn}(K \pm iK') = 0. \end{cases}$$

If in eqs. (14) we put $u^* = 0$, we see that as u approaches zero, the expressions

$$\operatorname{sn}(\pm iK'), \quad \operatorname{cn}(\pm iK'), \quad \operatorname{dn}(\pm iK')$$

approach infinity.

We see from what has preceded that Elliptic Functions have two periods, one a real period, and one an imaginary period.

In the former characteristic they resemble Trigonometric Functions, and in the latter Logarithmic Functions.

On account of these two periods they are often called Doubly Periodic Functions. Some authors make this double periodicity the starting-point of their investigations. This method of investigation gives some very beautiful results and processes, but not of a kind adapted for an elementary work.


It will be noticed that the Elliptic Functions $\operatorname{sn} u$, $\operatorname{cn} u$, and $\operatorname{dn} u$ have a very close analogy to trigonometric functions, in which, however, the independent variable u is not an angle, as it is in the case of trigonometric functions.

Like Trigonometric Functions, these Elliptic Functions can be arranged in tables. These tables, however, require a double argument, viz., u and k . In Chap. IX these functions are developed into series, from which their values may be computed and tables formed.

No complete tables have yet been published, though they are in process of computation.

CHAPTER IV.

LANDEN'S TRANSFORMATION.



LET AB be the diameter of a circle, with the centre at O , the radius $AO = r$, and C a fixed point situated upon OB , and $OC = k_0 r$. Denote the angle PBC by ϕ , and the angle PCO by ϕ_1 . Let P' be a point indefinitely near to P .

Then

$$\frac{PP'}{PC} = \frac{\sin PCP'}{\sin PP'C} = \frac{\sin PCP'}{\cos OP'C}$$

But $PP' = 2rd\phi$, and $\sin PCP' = PCP' = d\phi_1$;
therefore

$$\frac{2rd\phi}{PC} = \frac{d\phi_1}{\cos OP'C}$$

But

$$\begin{aligned} \overline{PC}^2 &= r^2 + r^2 k_0^2 + 2r^2 k_0 \cos 2\phi \\ &= (r + rk_0)^2 \cos^2 \phi + (r - rk_0)^2 \sin^2 \phi; \end{aligned}$$

also

$$\begin{aligned} r^2 \cos^2 OP'C &= r^2 - r^2 \sin^2 OP'C \\ &= r^2 - r^2 k_0^2 \sin^2 \phi_1. \end{aligned}$$

Therefore

$$\frac{2d\phi}{\sqrt{(r + rk_0)^2 \cos^2 \phi + (r - rk_0)^2 \sin^2 \phi}} = \frac{d\phi_1}{\sqrt{r^2 - r^2 k_0^2 \sin^2 \phi_1}},$$

which can be written

$$\frac{2}{r + rk_0} \frac{d\phi}{\sqrt{1 - \frac{4k_0^2 r^2}{(r + rk_0)^2} \sin^2 \phi}} = \frac{1}{r} \frac{d\phi_1}{\sqrt{1 - k_0^2 \sin^2 \phi_1}},$$

Putting

$$(1) \quad \frac{4k_0 r^2}{(r + rk_0)^2} = \frac{4k_0}{(1 + k_0)^2} = k^2,$$

we have

$$(2) \quad \int_0^\phi \frac{d\phi}{\sqrt{1 - k'^2 \sin^2 \phi}} = \frac{1 + k_0}{2} \int_0^{\phi_1} \frac{d\phi_1}{\sqrt{1 - k_0'^2 \sin^2 \phi_1}},$$

no constant being added because ϕ and ϕ_1 vanish simultaneously; ϕ and ϕ_1 being connected by the equation

$$(3) \quad \frac{\sin OPC}{\sin OCP} = \frac{\sin (2\phi - \phi_1)}{\sin \phi_1} = \frac{rk_0}{r} = k_0.$$

From the value of k^2 we have

$$(4) \quad 1 - k^2 = k'^2 = \frac{(1 - k_0)^2}{(1 + k_0)^2},$$

and therefore

$$(5) \quad k_0 = \frac{1 - k'}{1 + k'}.$$

k' is called the *complementary modulus*, and is evidently the minimum value of $\Delta\phi$, the value of $\Delta\phi$ when $\phi = 90^\circ$:

$$\sqrt{1 - k^2} = k'.$$

From eq. (1) we evidently have $k > k_0$, for, putting eq. (1) in the form

$$\frac{k^2}{k_0^2} = \frac{4}{k_0 + 2k_0^2 + k_0^3},$$

we see that if $k_0 = 1$, then $k = k_0$, but as $k_0 < 1$, always, as is evident from the figure, k must be greater than k_0 .

It is also evident, from the figure, that $\phi_1 > \phi$. Or it may be deduced directly from eq. (3).

Since $k < 1$, we can write

$$k = \sin \theta, \quad k' = \sqrt{1 - k^2} = \cos \theta.$$

Substituting in eq. (5), we have

$$k_0 = \frac{1 - k'}{1 + k'} = \tan^2 \frac{1}{2}\theta,$$

and we can write

$$k_0 = \sin \theta_0, \quad k_1' = \sqrt{1 - k_0^2} = \cos \theta_0.$$

From eq. (5) we have

$$1 + k_0 = \frac{2}{1 + k'}.$$

Substituting the value of k_0 in that for k_1' , we get

$$k_1' = \frac{2\sqrt{k'}}{1 + k'}.$$

We also have

$$2\phi - \phi_1 = \phi - (\phi_1 - \phi)$$

$$\phi_1 = \phi + (\phi_1 - \phi),$$

and, eq. (3),

$$\operatorname{sn} (2\phi - \phi_1) = k_0 \sin \phi_1,$$

becomes

$$\sin \varphi \cos (\phi_1 - \phi) - \cos \phi \sin (\phi_1 - \phi) =$$

$$k_0 \sin \phi \cos (\phi_1 - \phi) + k_0 \cos \phi \sin (\phi_1 - \phi),$$

or

$$\tan \phi - \tan (\phi_1 - \phi) = k_0 \tan \phi + k_0 \tan (\phi_1 - \phi),$$

or

$$\tan (\phi_1 - \phi) = \frac{1 - k_0}{1 + k_0} \tan \phi$$

$$= k' \tan \phi.$$

Collecting these results, we have

$$(6) \quad k = \frac{2\sqrt{k_0}}{1 + k_0} = \sin \theta;$$

$$(7) \quad k_0 = \frac{1 - k'}{1 + k'} = \sin \theta_0 = \tan^2 \frac{1}{2}\theta;$$

$$(8) \quad k_1' = \frac{2\sqrt{k'}}{1+k'} = \cos \theta_0;$$

$$(9) \quad k' = \frac{1-k_0}{1+k_0} = \cos \theta;$$

$$(10) \quad 1+k_0 = \frac{2}{1+k'} = \frac{2\sqrt{k_0}}{k} = \frac{k_1'}{\sqrt{k'}} = \frac{1}{\cos^2 \frac{1}{2}\theta};$$

$$(11) \quad \sin (2\phi - \phi_1) = k_0 \sin \phi_1;$$

$$(12) \quad \tan (\phi_1 - \phi) = k' \tan \phi;$$

$$(13) \quad \int_0^\phi \frac{d\phi}{\Delta(k, \phi)} = \frac{1+k_0}{2} \int_0^{\phi_1} \frac{d\phi_1}{\Delta(k_0, \phi_1)};$$

$$(14) \quad k = \sqrt{1-k'^2}, \quad k' = \sqrt{1-k^2}.$$

Upon examination it will easily appear that k and k_0 , and θ and θ_0 , are the first two terms of a decreasing series of moduli and angles; k' and k_1' , and ϕ and ϕ_1 , of an increasing series; the law connecting the different terms of the series being deduced from eqs. (6) to (12).

By repeated applications of these equations we would get the following series of moduli and amplitudes:

$k_{0n} = O_{(n=\infty)}$	$k_n' = I_{(n=\infty)}$	ϕ_n
$\vdots$	$\vdots$	$\vdots$
k_{00}	k_2'	ϕ_2
k_0	k_1'	ϕ_1
k	k'	ϕ

The upper limit of the one series of moduli is 1, and the lower limit of the other series is 0, as is indicated. k and k' ,

which are bound by the relation $k^2 + k'^2 = 1$, are called the *primitives* of the series.

NOTE.—It will be noticed that the successive terms of a decreasing series are indicated by the sub-accents 0, 00, 03, 04, . . . 0*n*; and the successive terms of an increasing series by the sub-accents 1, 2, 3, . . . *n*.

Again, by application of these equations, we can form a new series running up from *k*, viz., $k_1, k_2, k_3, \dots k_n = 1_{(n=\infty)}$; and also a new series running down from k' , viz., $k'_0, k'_{00}, \dots k'_{0n} = 0_{(n=\infty)}$. So also with ϕ .

Collecting these series, we have

$k_{0n} = 0$	$k'_n = 1$	ϕ_n
⋮	⋮	⋮
⋮	⋮	⋮
⋮	⋮	⋮
k_{03}	k'_2	ϕ_2
k_0	k'_1	ϕ_1
k	k'	ϕ
k_1	k'_0	ϕ_0
k_2	k'_{00}	ϕ_{00}
⋮	⋮	⋮
⋮	⋮	⋮
⋮	⋮	⋮
$k_n = 1$	$k'_{0n} = 0$	$\phi_{0n} = 0$

NOTE.—In practice it will be found that generally *n* will not need to be very large in order to reach the limiting values of the terms, often only two or three terms being needed.

Applying eqs. (7), (12), (13), and (14) repeatedly, we get

$$(14) \left\{ \begin{array}{ll} k = \sin \theta, & k' = \cos \theta; \\ k_0 = \frac{1 - k'}{1 + k'} = \tan^2 \frac{1}{2} \theta = \sin \theta_0, & k'_1 = \cos \theta_0; \\ k_{00} = \tan^2 \frac{1}{2} \theta_0 = \sin \theta_{00}, & k'_2 = \cos \theta_{00}; \\ k_{03} = \tan^2 \frac{1}{2} \theta_{00} = \sin \theta_{03}, & k'_3 = \cos \theta_{03}; \\ \dots & \dots \\ k_{0n} = \tan^2 \frac{1}{2} \theta_{0(n-1)} = \sin \theta_{0n}, & k'_n = \cos \theta_{0n}. \end{array} \right.$$

$$(14_1) \quad \left\{ \begin{array}{l} \tan (\phi_1 - \phi) = k' \tan \phi; \\ \tan (\phi_2 - \phi_1) = k'_1 \tan \phi_1; \\ \tan (\phi_3 - \phi_2) = k'_2 \tan \phi_2; \\ . \quad . \quad . \quad . \quad . \quad . \quad . \\ \tan (\phi_n - \phi_{n-1}) = k'^{(n-1)} \tan \phi_{n-1}. \end{array} \right.$$

From eqs. (15) and (10) we get

$$F(k, \phi) = \sqrt{\frac{k'_1 k'_2 k'_3 \dots k'_n}{k'}} \cdot \frac{F(k_{0n}, \phi_n)}{2^n}.$$

And this with equations (8) and (9) gives

$$(17) \quad F(k, \phi) = \sqrt{\frac{\cos \theta_0 \cos \theta_{00} \dots \cos^2 \theta_{0n}}{\cos \theta}} \cdot \frac{F(k_{0n}, \phi_n)}{2^n}.$$

Applying equation (13) to (k_1, ϕ_0) , (k_2, ϕ_{00}) , etc., we get

$$F(k_1, \phi_0) = \frac{1+k}{2} F(k, \phi), \text{ etc.};$$

but since, eq. (10),

$$\frac{1+k}{2} = \frac{1}{1+k'_0}, \text{ etc.},$$

these become

$$F(k, \phi) = (1+k'_0)F(k_1, \phi_0);$$

$$F(k_1, \phi_0) = (1+k'_{00})F(k_2, \phi_{00});$$

$$\dots \dots \dots$$

$$F(k_{n-1}, \phi_{0(n-1)}) = (1+k'_{0n})F(k_{0n}, \phi_{0n});$$

whence

$$(18) \quad F(k, \phi) = (1+k'_0)(1+k'_{00}) \dots (1+k'_{0n})F(k_{0n}, \phi_{0n}),$$

in which k'_0 , k'_{00} , etc., k_1 , k_2 , etc., ϕ_0 , ϕ_{00} , etc., are determined as follows:

$$\begin{aligned} \text{Let} \quad k &= \sin \theta, \\ k_1 &= \sin \theta_1. \end{aligned}$$

From eq. (10),

$$k_1 = \frac{2\sqrt{k}}{1+k} \quad \text{or} \quad \sin \theta_1 = \frac{2\sqrt{\sin \theta}}{1+\sin \theta}.$$

Solving this equation for $\sin \theta$, we get

$$\sin \theta = \tan^2 \frac{1}{2} \theta_1.$$

Hence we can write

$$(18_1) \quad \left\{ \begin{array}{l} k = \sin \theta = \tan^2 \frac{1}{2} \theta_1; \\ k_1 = \sin \theta_1 = \tan^2 \frac{1}{2} \theta_2; \\ \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \\ k_n = \sin \theta_n. \end{array} \right.$$

From equation (12) we get

$$(18_2) \quad \left\{ \begin{array}{l} \sin (2\phi_0 - \phi) = k \sin \phi; * \\ \sin (2\phi_{00} - \phi_0) = k_1 \sin \phi_0; \\ \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \\ \sin (2\phi_{0n} - \phi_{0(n-1)}) = k_{n-1} \sin \phi_{0(n-1)}. \end{array} \right.$$

* When $\sin \phi = 1$ nearly, ϕ is best determined as follows: From eq. (12) we have

$$\begin{aligned} \tan (\phi - \phi_0) &= k'_0 \tan \phi_0 \\ &= k'_0 \tan \phi \text{ nearly;} \end{aligned}$$

whence

$$\phi - \phi_0 = R k'_0 \tan \phi \text{ nearly,}$$

R being the radian in seconds, viz. 206264''.806, and $\log R = 5.3144251$.

Substituting the approximate value of ϕ_0 , we can get a new approximation.

Example. $\phi_0 = 82^\circ 30'$ $k'_{00} = \log^{-1} 5.8757219$

$\tan 82^\circ 30'$	10.8805709	
k'_{00}	5.8757219	
R	5.3144251	
	2.0707179	$117''.684 = 1'.9614$

$$\phi_0 - \phi_{00} = 1'.9614$$

$$\phi_{00} = 82^\circ 28'.0386$$

1st approximation.

To determine k'_0 , k'_{00} , etc., we have

$$(18_s) \quad \begin{cases} k' = \sin \eta, & k = \cos \eta; \\ k'_0 = \frac{1-k}{1+k} = \tan^2 \frac{1}{2}\eta = \sin \eta_0, & k_1 = \cos \eta_0; \\ k'_{00} = \tan^2 \frac{1}{2}\eta_0 = \sin \eta_{00}, & k_2 = \cos \eta_{00}; \\ \text{etc.} & \text{etc.} & \text{etc.} \end{cases}$$

Or, since $1 + k'_0 = \frac{1}{\cos^2 \frac{1}{2}\eta}$, $1 + k'_{00} = \frac{1}{\cos^2 \frac{1}{2}\eta_0}$, etc.,

we can put eq. (18) in the following form :

$$(19) \quad F(k, \phi) = \frac{1}{\cos^2 \frac{1}{2}\eta \cos^2 \frac{1}{2}\eta_0 \dots \cos^2 \frac{1}{2}\eta_{0n}} F(k_n, \phi_{0n}).$$

From equation (13) we have

$$(19)^* \quad F(k_1, \phi_0) = \frac{1+k}{2} F(k, \phi),$$

whence

$$F(k, \phi) = \frac{2}{1+k} F(k_1, \phi_0).$$

By repeated applications this gives, after combining,

$$F(k, \phi) = \frac{2}{1+k} \cdot \frac{2}{1+k_1} \dots \frac{2}{1+k_{n-1}} \cdot F(k_n, \phi_{0n})$$

This value gives

$$\phi_0 - \phi_{00} = 117''.1675 = 1'.95279$$

$$\therefore \phi_{00} = 82^\circ 28'.04721$$

2d approximation.

This value gives

$$\phi_0 - \phi_{00} = 117''.1698 = 1'.95283$$

$$\phi_{00} = 82^\circ 28'.04717$$

3d approximation.

$$= \frac{k_1}{\sqrt{k}} \cdot \frac{k_2}{\sqrt{k_1}} \cdots \frac{k_n}{\sqrt{k_{n-1}}} \cdot F(k_n, \phi_{0n});$$

$$(20) \quad F(k, \phi) = \sqrt{\frac{k_1 k_2 \cdots k_n^2}{k}} \cdot F(k_n, \phi_{0n});$$

k_1, k_2 , etc., being determined by repeated applications of

$$k_1 = \frac{2\sqrt{k}}{1+k},$$

or by equations (18₁).

In equation (19)* let us change k_1 and ϕ_0 into k' and ϕ respectively, so that the first member may have for its complete function

$$K' = F(k', \phi).$$

Upon examination of eq. (19)* we see that the modulus in the second member must be the one next less than the one in the first member, that is, k'_0 ; and likewise that the amplitude must be the one next greater than the amplitude in the first member, viz., ϕ_1 ; hence we get

$$F(k', \phi) = \frac{1+k'_0}{2} F(k'_0, \phi_1).$$

Indicating the complete functions by K' and K'_0 , we have, since $\phi = \frac{\pi}{2}$ when $\phi_1 = \pi$ (see Chap. V),

$$K' = (1 + k'_0)K'_0;$$

and in the same manner,

$$\begin{aligned} K'_0 &= (1 + k'_{00})K'_{00}, \\ K'_{00} &= (1 + k'_{03})K'_{03}, \\ &\cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \\ K'_{0(n-1)} &= (1 + k'_{0n})K'_{0n}; \end{aligned}$$

whence

$$K' = (1 + k'_0)(1 + k'_{00}) \dots (1 + k'_{0n})K'_{0n}.$$

Since

$$K'_{0n} = \int_0^{\frac{\pi}{2}} d\phi = \frac{\pi}{2}, \quad (n = \text{limit}),$$

we have

$$(20)^* \quad (1 + k'_0)(1 + k'_{00}) \dots (1 + k'_{0n}) = \frac{2K'}{\pi}.$$

From eq. (19)* we have, since [eq. (10), Chap IV]

$$\frac{1 + k}{2} = \frac{1}{1 + k'_0},$$

$$(1 + k'_0) \int_0^{\phi_0} \frac{d\phi_0}{\Delta(\phi_0, k_1)} = \int_0^{\phi} \frac{d\phi}{\Delta(\phi_1, k)};$$

whence also, since for $\phi_0 = \frac{\pi}{2}$, $\phi = \pi$,

$$\begin{aligned} (1 + k'_0) K_1 &= 2K, \\ (1 + k'_{00}) K_2 &= 2K_1, \\ &\dots \dots \dots \\ (1 + k'_{0n}) K_n &= 2K_{n-1}, \end{aligned}$$

and

$$(1 + k'_0)(1 + k'_{00}) \dots (1 + k'_{0n})K_n = 2^n K;$$

or

$$\frac{K_n}{2^n} = \frac{K}{(1 + k'_0)(1 + k'_{00}) \dots} \quad (n = \infty)$$

$$(21) \quad = \frac{\pi}{2K_1} K.$$

Let us find the limiting value of $F(k_{0n}, \phi_n)$ in eq. (15). In the equation $\tan(\phi_n - \phi_{n-1}) = k_{n-1} \tan \phi_{n-1}$, we see that when k_{n-1} reaches the limit 1, then $\phi_n - \phi_{n-1} = \phi_{n-1}$ or $\phi_n = 2\phi_{n-1}$. Therefore

$$\frac{\phi_n}{2^n} = \frac{2\phi_{n-1}}{2^n} = \frac{\phi_{n-1}}{2^{n-1}};$$

$$\frac{\phi_{n+1}}{2^{n+1}} = \frac{2\phi_n}{2^{n+1}} = \frac{\phi_n}{2^n} = \frac{\phi_{n-1}}{2^{n-1}};$$

$$\frac{\phi_{n+m}}{2^{n+m}} = \frac{\phi_{n-1}}{2^n} = \text{constant, whatever } m \text{ may be.}$$

Therefore eq. (15) becomes

$$(21)^* \quad F(k, \phi) = (1 + k_0)(1 + k_{00}) \dots (1 + k_{0n}) \frac{\phi_n}{2^n},$$

n being whatever number will carry k_0 and $\frac{\phi_1}{2}$ to their limiting values.

In the same way, eqs. (16) and (17) become

$$(22) \quad F(k, \phi) = \frac{1}{\cos^2 \frac{\theta}{2} \cos^2 \frac{\theta_0}{2} \dots \cos^2 \frac{\theta_{0n}}{2}} \cdot \frac{\phi_n}{2^n}$$

$$(23) \quad = \sqrt{\frac{\cos \theta_0 \cos \theta_{00} \dots \cos^2 \theta_{0n}}{\cos \theta}} \cdot \frac{\phi_n}{2^n},$$

$n - 1$ being the number which makes $k'_{n-1} = 1$.

In these last three equations k_0, k_{00} are determined by eqs. (14₁); ϕ_1, ϕ_2 , etc., by eqs. (14₂)*; θ, θ_0 , etc., by eqs. (14₁); and k', k'_0, k'_2 , etc., for use in eq. (14₂) by eqs. (14₁).

* Taking for $\phi_1 - \phi$, etc., not always the least angle given by the tables, but that which is nearest to ϕ .

BISECTED AMPLITUDES.

We have identically

$$u = 2 \cdot \frac{u}{2} = 2 \int \frac{d \operatorname{am} \frac{u}{2}}{\sqrt{1 - k^2 \operatorname{sn}^2 \frac{u}{2}}};$$

$$\frac{u}{2} = 2 \cdot \frac{u}{4} = 2F\left(k, \operatorname{am} \frac{u}{4}\right);$$

etc.

Therefore

$$\begin{aligned} u &= F(k, \operatorname{am} u) = 2^n F\left(k, \operatorname{am} \frac{u}{n}\right) \\ &= 2^n \cdot \operatorname{am} \frac{u}{n}, \quad (n = \text{limit}), \end{aligned}$$

$\operatorname{am} \frac{u}{n}$ being determined by repeated applications of eq. (12) of Chap. II, as follows:

$$\operatorname{sn}^2 \frac{u}{2} = \frac{1 - \operatorname{cn} u}{1 + \operatorname{dn} u} = \frac{2 \sin^2 \frac{1}{2} \operatorname{am} u}{1 + \operatorname{dn} u};$$

$$(24) \quad \operatorname{sn} \frac{u}{2} = \frac{\sin \frac{1}{2} \operatorname{am} u}{\sqrt{\frac{1 + \cos \beta}{2}}} = \frac{\sin \frac{\operatorname{am} u}{2}}{\cos \frac{1}{2} \beta};$$

β being an angle determined by the equation

$$(25) \quad \cos \beta = \operatorname{dn} u = \sqrt{1 - k^2 \operatorname{sn}^2 u},$$

and n being the number which makes

$$2^n \operatorname{am} \frac{u}{n} = \text{constant}.$$

$\operatorname{am} \frac{u}{n}$ is found by repeated applications of eq. (24).

Indicating the amplitudes as follows :

$$\text{am } u = \phi,$$

$$\text{am } \frac{u}{2} = \phi_{02},$$

$$\text{am } \frac{u}{4} = \phi_{04},$$

$$\text{am } \frac{u}{8} = \phi_{08},$$

$$\dots$$

$$\text{am } \frac{u}{2^n} = \phi_{02^n},$$

$$(26) \quad F(k, \phi) = 2^n \phi_{02^n};$$

n being the limiting value.

In eq. (18), when k_n reaches its limit 1, we have

$$F(k_n, \phi_{0n}) = \int_0^\phi \frac{d\phi_{0n}}{\cos \phi_{0n}} = \log_e \tan (45^\circ + \tfrac{1}{2}\phi_{0n}),$$

and eqs. (18) and (19) become

$$(27) \quad F(k, \phi) = (1 + k'_0)(1 + k'_{00}) \dots (1 + k'_{0n}) \log_e \tan (45^\circ + \tfrac{1}{2}\phi_{0n})$$

$$= \frac{1}{\cos^2 \tfrac{1}{2}\eta \cos^2 \tfrac{1}{2}\eta_0 \dots \cos^2 \tfrac{1}{2}\eta_{0n}} \log_e \tan (45^\circ + \tfrac{1}{2}\phi_{0n})$$

$$(28) \quad = \frac{1}{\cos^2 \tfrac{1}{2}\eta \cos^2 \tfrac{1}{2}\eta_0 \dots \cos^2 \tfrac{1}{2}\eta_{0n}} \cdot \frac{1}{M} \log \tan (45^\circ + \tfrac{1}{2}\phi_{0n});$$

n being the number which renders $k_n = 1$.

Eq. (20) becomes

$$\begin{aligned}
 (29) \quad F(k, \phi) &= \sqrt{\frac{k_1 k_2 \dots k_n}{k}} \cdot \log_e \tan (45^\circ + \tfrac{1}{2} \phi_{0n}) \\
 &= \sqrt{\frac{k_1 k_2 \dots k_n}{k}} \cdot \frac{1}{M} \log \tan (45^\circ + \tfrac{1}{2} \phi_{0n}) \\
 &= \sqrt{\frac{\cos \eta_0 \cos \eta_{00} \dots \cos^2 \eta_{0n}}{\cos \eta}} \cdot \frac{1}{M} \log \tan (45^\circ + \tfrac{1}{2} \phi_{0n}).
 \end{aligned}$$

In these equations k'_0, k'_{00} , etc., are determined by eqs. (18₃); η, η_0 , etc., by eqs. (18₂); ϕ_0, ϕ_{00} , etc., by eqs. (18₂); k_1, k_2 , etc., by eqs. (18₁).

Substituting in eq. (27) from eq. (20)*, we have

$$\begin{aligned}
 F(k, \phi) &= \frac{2K'}{\pi} \log_e \tan (45^\circ + \tfrac{1}{2} \phi_{0n}) \\
 (30) \quad &= \frac{2K'}{\pi M} \log \tan (45^\circ + \tfrac{1}{2} \phi_{0n}).
 \end{aligned}$$

CHAPTER V.

COMPLETE FUNCTIONS.

INDICATE by K the complete integral

$$(1) \quad K = \int_0^{\frac{\pi}{2}} \frac{d\phi}{\sqrt{1 - k^2 \sin^2 \phi}},$$

and by K_0 the complete integral

$$(2) \quad K_0 = \int_0^{\frac{\pi}{2}} \frac{d\phi_1}{\sqrt{1 - k_0^2 \sin^2 \phi_1}};$$

and in a similar manner K_{00} , K_{03} , etc.

From eq. (12), Chap. IV, we have

$$\begin{aligned} \tan(\phi_1 - \phi) &= k' \tan \phi \\ &= \frac{\tan \phi_1 - \tan \phi}{1 + \tan \phi_1 \tan \phi}, \end{aligned}$$

whence

$$\begin{aligned} \tan \phi_1 &= \frac{(1 + k') \tan \phi}{1 - k' \tan^2 \phi} \\ &= \frac{1 + k'}{\frac{1}{\tan \phi} - k' \tan \phi}. \end{aligned}$$

From this equation we see that when $\phi = \frac{\pi}{2}$, $\phi_1 = \pi$. This same result might also have been deduced from Fig. 1, Chap. IV, or from the equation

$$(3) \quad \phi_1 = 2\phi - k_0 \sin 2\phi + \frac{1}{2}k_0^2 \sin 4\phi - \text{etc.},$$

this last being the well-known trigonometrical formula

$$\tan x = n \tan y,$$

$$x = y - \frac{1-n}{1+n} \sin 2y + \frac{1}{2} \left(\frac{1-n}{1+n} \right)^2 \sin 4y - \frac{1}{3} \left(\frac{1-n}{1+n} \right)^3 \sin 6y + \text{etc.}$$

Since $\int_0^{\frac{\pi}{2}} \frac{d\phi_1}{\Delta(k_0\phi_1)} = K_0$, we must have

$$\int_0^{\pi} \frac{d\phi_1}{\Delta(k_0\phi_1)} = 2K_0.$$

These values substituted in eq. (13), Chap. IV, give successively

$$(4) \quad \begin{aligned} K &= (1 + k_0)K_0, \\ K_0 &= (1 + k_{00})K_{00}, \\ &\dots \dots \dots \\ K_{0(n-1)} &= (1 + k_{0n})K_{0n}; \end{aligned}$$

whence

$$(5) \quad K = (1 + k_0)(1 + k_{00}) \dots (1 + k_{0n})K_{0n}.$$

Since the limit of k_{0n} is 0, K_{0n} becomes

$$K_{0n} = \int_0^{\frac{\pi}{2}} d\phi = \frac{\pi}{2},$$

and we have

$$(6) \quad K = \frac{\pi}{2} (1 + k_0)(1 + k_{00}) \dots$$

$$(7) \quad = \frac{\frac{1}{2}\pi}{\cos^2 \frac{1}{2}\theta \cos^2 \frac{1}{2}\theta_0 \dots \cos^2 \frac{1}{2}\theta_{0n}};$$

k_1, k_0 , etc., and θ_1, θ_0 , etc., being found by eqs. (14₁) of Chap. IV.

From the formulæ in these two chapters we can compute the values of u for all values of ϕ and k and arrange them in tables. These are Legendre's Tables of Elliptic Integrals.

CHAPTER VI.

EVALUATION FOR ϕ .

TO FIND ϕ , u AND k BEING GIVEN.

From eqs. (21) and (23), Chap. IV, we have (n having the value which makes $\cos \theta_{0n} = 1$)

$$(1) \quad \phi_n = \frac{2^n u}{(1 + k_0)(1 + k_{00}) \dots (1 + k_{0n})} = \frac{2^n u \sqrt{\cos \theta}}{\sqrt{\cos \theta_0 \dots \cos^2 \theta_{0n}}},$$

from which ϕ_n can be calculated, k_0, k_{00} , etc., being found by means of equations (14₁), Chap. IV.

Then, having ϕ_n, k_0, k_{00} , etc., we can find ϕ by means of the following equations:

$$\begin{aligned} \sin (2\phi_{n-1} - \phi_n) &= k_{0n} \sin \phi_n, \\ \sin (2\phi_{n-2} - \phi_{n-1}) &= k_{0(n-1)} \sin \phi_{n-1}, \\ &\dots \dots \dots \\ \sin (2\phi - \phi_1) &= k_0 \sin \phi_1; \end{aligned}$$

whence we can get the angle ϕ .

When $k > \sqrt{\frac{1}{2}}$ the following formulæ will generally be found to work more rapidly:

From eq. (29), Chap. IV, we have

$$(2) \quad \log \tan (45^\circ + \tfrac{1}{2}\phi_{0n}) = \frac{uM}{\sqrt{\frac{k_1 k_2 \dots k_n^2}{k}}},$$

from which we can get ϕ_{0n} ; k_1, k_2 , etc., being calculated from eqs. (18), Chap. IV, and ϕ being calculated from the following equations:

$$\tan (\phi_{0(n-1)} - \phi_{0n}) = k_n \tan \phi_{0n},$$

$$\dots \dots \dots$$

$$\tan (\phi_0 - \phi_{00}) = k_0 \tan \phi_{00},$$

$$\tan (\phi - \phi_0) = k \tan \phi_0;$$

whence we get ϕ .

This gives a method of solving the equation

$$F\psi = nF\phi,$$

where n and ϕ and the moduli are known, and ψ is the required quantity. n and ϕ give $F\psi$, and then ψ can be determined by the foregoing methods.

When $k = 1$ nearly, equation (2) takes a special form,—

1°. When $\tan \phi$ is very much less than $\frac{1}{k'}$. In this case

$$\begin{aligned} F(k, \phi) &= \int \frac{d\phi}{\sqrt{\cos^2 \phi + k'^2 \sin^2 \phi}} = \int \frac{d\phi}{\sqrt{(1 + k'^2 \tan^2 \phi) \cos^2 \phi}} \\ &= \int \frac{d\phi}{\cos \phi} = \log \tan (45^\circ + \tfrac{1}{2}\phi); \end{aligned}$$

whence we can find ϕ .

2°. When $\tan \phi$ and $\frac{1}{k'}$ approach somewhat the same value, and $k' \tan \phi$ cannot be neglected, $F(k, \phi)$ must be transposed into another where k' shall be much smaller, so that $k' \tan \phi$ can be neglected.

These methods for finding ϕ apply only when $\phi < \frac{\pi}{2}$, that is, $u < K$. In the opposite case ($u > K$) put

$$u = 2nK \pm v,$$

the upper or the lower sign being taken according as K is continued in u an even or an odd number of times. In either case $v < K$, and we can find v by the preceding methods.

Having found v , we have from eq. (5), Chap. III,

$$\begin{aligned} \text{am } u &= \text{am } (2nK \pm v) \\ &= n\pi \pm \text{am } v. \end{aligned}$$

CHAPTER VII.

DEVELOPMENT OF ELLIPTIC FUNCTIONS INTO FACTORS.

FROM eq. (12), Chap. IV, we readily get

$$\sin (2\phi_0 - \phi) = k \sin \phi;$$

$$\begin{aligned} \sin \phi &= \frac{\sin 2\phi_0}{\sqrt{1 + k^2 + 2k \cos 2\phi_0}} \\ &= \frac{\sin 2\phi_0}{\sqrt{(1 + k^2) - 4k \sin^2 \phi_0}} \\ &= \frac{1 + k_0'}{2} \cdot \frac{\sin 2\phi_0}{\sqrt{1 - k_1^2 \sin^2 \phi_0}} \end{aligned}$$

(since $\frac{4k}{1+k} = k_1$ and $1+k = \frac{2}{1+k_0'}$, eqs. (6) and (10), Chap. IV);
and thence

$$(1) \quad \sin \phi = \frac{(1 + k_0') \sin \phi_0 \cos \phi_0}{\Delta(\phi_0, k_1)}.$$

From eq. (13), Chap. IV, we have

$$\int_0^{\phi_0} \frac{d\phi_0}{\Delta(\phi_0, k_1)} = \frac{1 + k}{2} \int_0^{\phi} \frac{d\phi}{\Delta(\phi, k)};$$

and from eq. (4), Chap. V, passing up the scale of moduli one step,

$$1 + k = \frac{K_1}{K},$$

whence

$$F(\phi_0, k_1) = \frac{K_1}{2K} F(\phi, k).$$

Put

$$F(\phi_0, k_1) = u_1 \quad \text{and} \quad F(\phi, k) = u,$$

whence

$$u_1 = \frac{K_1}{2K} u.$$

Furthermore,

$$\phi = \text{am}(u, k);$$

$$\phi_1 = \text{am}(u_1, k_1) = \text{am}\left(\frac{K_1}{2K} u, k_1\right).$$

Substituting these values in eq. (I), we have

$$\text{sn}(u, k) = (1 + k_0') \frac{\text{sn}\left(\frac{K_1}{2K} u, k_1\right) \text{cn}\left(\frac{K_1}{2K} u, k_1\right)}{\text{dn}\left(\frac{K_1}{2K} u, k_1\right)}.$$

But from eq. (II), Chap. III, we have

$$\frac{\text{cn}(\nu, k_1)}{\text{dn}(\nu, k_1)} = \text{sn}(\nu + K_1, k_1),$$

or

$$\begin{aligned} \frac{\text{cn}\left(\frac{K_1}{2K} u, k_1\right)}{\text{dn}\left(\frac{K_1}{2K} u, k_1\right)} &= \text{sn}\left(\frac{K_1 u}{2K} + K_1, k_1\right) \\ &= \text{sn}\left(\frac{K_1}{2K} (u + K_1), k_1\right); \end{aligned}$$

whence

$$(2) \quad \operatorname{sn}(u, k) = (1 + k'_0) \operatorname{sn} \frac{K_1 u}{2K} \operatorname{sn} \left[\frac{K_1}{2K} (u + 2K) \right]^\dagger$$

(Mod. = k_1 .)

From this equation, evidently, we have generally

$$(2)^* \quad \operatorname{sn}(\nu, k_n) = (1 + k'_{0(n+1)}) \operatorname{sn} \frac{K_{n+1}}{2K_n} \nu \operatorname{sn} \left[\frac{K_{n+1}}{2K_n} (\nu + 2K_n) \right].$$

(Mod. = k_{n+1} .)

Applying this general formula to the two factors of eq. (2), we have

$$\operatorname{sn} \left(\frac{K_1 u}{2K}, k_1 \right) = (1 + k'_{00}) \operatorname{sn} \frac{K_2}{2K_1} \cdot \frac{K_1 u}{2K} \cdot \operatorname{sn} \left[\frac{K_2}{2K_1} \left(\frac{K_1 u}{2K} + 2K_1 \right) \right]$$

(Mod. k_2)

$$= (1 + k'_{00}) \operatorname{sn} \frac{K_2 u}{2^2 K} \operatorname{sn} \frac{K_2}{2^2 K} (u + 4K); \quad (\text{Mod. } k_2);$$

$$(3) \quad \operatorname{sn} \left[\frac{K_1}{2K} (u + 2K), k_1 \right] = (1 + k'_{00}) \operatorname{sn} \frac{K_2}{2^2 K} (u + 2K)$$

$$\cdot \operatorname{sn} \frac{K_2}{2K_1} \left[\frac{K_1}{2K} (u + 2K) + 2K_1 \right]. \quad (\text{Mod. } k_2.)$$

The last argument in this equation is equal to

$$\frac{K_2}{2^2 K} (u + 6K);$$

and since, eq. (7), Chap. III,

$$\operatorname{sn}(u, k_2) = \operatorname{sn}(2K_2 - u, k_2),$$

† The analogous formula in Trigonometry is

$$\sin \phi = \frac{1}{2} \sin \frac{1}{2} \phi \sin \frac{1}{2} (\phi + \pi).$$

we can put in place of this,

$$2K_2 - \frac{K_2}{2^2 K} (u + 6K = \frac{K_2}{2^2 K} (2K - u);$$

whence eq. (3) becomes

$$\begin{aligned} \operatorname{sn} \left[\frac{K_1}{2K} (u + 2K), k_1 \right] &= (1 + k'_{00}) \operatorname{sn} \frac{K_2}{2^2 K} (2K + u) \\ &\quad \cdot \operatorname{sn} \frac{K_2}{2^2 K} (2K - u). \end{aligned} \quad (\text{Mod. } k_2.)$$

Substituting these values in eq. (2), we have

$$\begin{aligned} (4) \quad \operatorname{sn} (u, k) &= (1 + k_0)(1 + k'_{00})^2 \operatorname{sn} \frac{K_2 u}{2^2 K} \\ &\quad \cdot \operatorname{sn} \frac{K_2}{2^2 K} (2K \pm u) \operatorname{sn} \frac{K_2}{2^2 K} (4K + u), \quad (\text{Mod. } k_2,) \end{aligned}$$

in which the double sign indicates two separate factors which are to be multiplied together.

By the application of the general equation (2)* we find that the arguments in the second member of eq. (4) will each give rise to two new arguments, as follows:

$$\frac{K_2 u}{2^2 K} \text{ gives } \frac{K_3 u}{2^2 K},$$

and

$$\frac{K_2}{2^2 K} \left(\frac{K_2 u}{2^2 K} + 2K \right) = \frac{K_3}{2^2 K} (u + 8K);$$

$$\frac{K_2}{2^2 K} (2K \pm u) \text{ gives } \frac{K_3}{2^2 K} (2K \pm u),$$

and

$$\frac{K_3}{2K_2} \left[\frac{K_2}{2^2 K} (2K \pm u) + 2K_2 \right] = \frac{K_3}{2^3 K} (10K \pm u), \quad . \quad . \quad (a)$$

$$\frac{K_3}{2^3 K} (4K + u) \quad \text{gives} \quad \frac{K_3}{2^3 K} (4K + u),$$

and

$$\frac{K_3}{2K_2} \left[\frac{K_2}{2^2 K} (4K + u) + 2K_2 \right] = \frac{K_3}{2^3 K} (12K + u). \quad . \quad . \quad (b)$$

Subtracting (a) and (b) from $2K_3$, by which the sine of the amplitudes will not be changed [eq. (7), Chap. III], and since our new modulus is k_3 , we have for the expressions (a) and (b),

$$\frac{K_3}{2^3 K} (6K \mp u); \quad . \quad . \quad . \quad . \quad . \quad (a')$$

$$\frac{K_3}{2^3 K} (4K - u). \quad . \quad . \quad . \quad . \quad . \quad (b')$$

Substituting these values in eq. (4), and remembering the factor $(1 + k'_{03})$ introduced by each application of eq. (2)*, we have

$$\text{sn } (u, k) = (1 + k'_0)(1 + k'_{00})^2(1 + k'_{03})^4 \text{sn } \frac{K_3 u}{2^3 K};$$

$$\cdot \text{sn } \frac{K_3}{2^3 K} (2K \pm u) \text{sn } \frac{K_3}{2^3 K} (4K \pm u);$$

$$\cdot \text{sn } \frac{K_3}{2^3 K} (6K \pm u) \text{sn } \frac{K_3}{2^3 K} (8K + u). \quad (\text{Mod. } k_3.)$$

From this the law governing the arguments is clear, and we can write for the general equation

$$\begin{aligned}
 (5) \quad \operatorname{sn}(u, k) &= (1 + k'_0)(1 + k'_{00})^2(1 + k'_{03})^4 \dots (1 + k'_{0n})^{2^{n-1}} \\
 &\cdot \operatorname{sn} \frac{K_n u}{2^n K} \operatorname{sn} \frac{K_n}{2^n K} (2K \pm u) \\
 &\cdot \operatorname{sn} \frac{K_n}{2^n K} (4K \pm u) \operatorname{sn} \frac{K_n}{2^n K} (6K \pm u) \\
 &\cdot \dots \operatorname{sn} \frac{K_n}{2^n K} \left[(2^n - 2)K \pm u \right] \\
 &\cdot \operatorname{sn} \frac{K_n}{2^n K} (2^n K + u). \quad (\text{Mod. } k_n)
 \end{aligned}$$

Indicate the continued product of the binomial factors by A' , and we have

$$A' = (1 + k'_0)(1 + k'_{00})^2(1 + k'_{03})^4(1 + k'_{04})^8 \dots$$

Since the limit of k'_0 , k'_{00} , etc., is zero, it is evident that these factors converge toward the value unity. It can be shown that the functional factors also converge toward the value unity. Thus the argument of the last factor can be written

$$K_n + \frac{K_n u}{2^n K}.$$

From eq. (11), Chap. III, we get then

$$\operatorname{sn} \left(K_n + \frac{K_n u}{2^n K} \right) = \frac{\operatorname{cn} \frac{K_n u}{2^n K}}{\operatorname{dn} \frac{K_n u}{2^n K}}. \quad (\text{Mod. } k_n)$$

But since k_n at its limit is equal to unity, $\operatorname{cn} = \operatorname{dn}$; whence the last factor of eq. (5) is unity.

From eq. (21), Chap. IV, we have

$$\lim_{n \rightarrow \infty} \frac{K_n}{2^n K'} = \frac{2\pi}{2K'}.$$

Therefore for $n = \infty$, eq. (5) becomes

$$\begin{aligned} \operatorname{sn}(u, k) = A' \operatorname{sn} \frac{\pi u}{2K'} \operatorname{sn} \frac{\pi}{2K'} (2K \pm u) \\ \cdot \operatorname{sn} \frac{\pi}{2K'} (4K \pm u) \operatorname{sn} \frac{\pi}{2K'} (6K \pm u), \quad \dots \dots \dots \\ (\text{Mod. } 1,) \end{aligned}$$

or

$$(6) \quad \operatorname{sn}(u, k) = A' \operatorname{sn} \frac{\pi u}{2K'} \left[\prod_h^\infty \right] \operatorname{sn} \frac{\pi}{2K'} (2hK \pm u),$$

(Mod. 1,)

where the sign $[II]$ indicates the continued product in the same manner as Σ indicates the continued sum.

When $k = 1$, $\int_0^\phi F(\phi, k)$ becomes

$$\nu = \int_0^\phi \frac{d\phi}{\cos \phi} = \frac{1}{2} \log \frac{1 + \sin \phi}{1 - \sin \phi};$$

whence

$$e^{2\nu} = \frac{1 + \sin \phi}{1 - \sin \phi}$$

and

$$\sin \phi = \frac{e^{2\nu} - 1}{e^{2\nu} + 1} = \frac{e^\nu - e^{-\nu}}{e^\nu + e^{-\nu}} = \operatorname{sn}(\nu, 1).$$

Hence in equation (6)

$$\sin \frac{\pi u}{2K'} = \frac{e^{\frac{\pi u}{2K'}} - e^{-\frac{\pi u}{2K'}}}{e^{\frac{\pi u}{2K'}} + e^{-\frac{\pi u}{2K'}}};$$

$$\operatorname{sn} \frac{\pi}{2K'} (2hK \pm u) = \frac{e^{\frac{h\pi K}{K'}} e^{\pm \frac{\pi u}{2K'}} - e^{-\frac{h\pi K}{K'}} e^{\mp \frac{\pi u}{2K'}}}{e^{\frac{h\pi K}{K'}} e^{\pm \frac{\pi u}{2K'}} + e^{-\frac{h\pi K}{K'}} e^{\mp \frac{\pi u}{2K'}}}.$$

Put

$$(6)^* \quad q = e^{-\frac{\pi K'}{K}}, \quad q' = e^{-\frac{\pi K}{K'}},$$

and the last expression becomes

$$\operatorname{sn} \frac{\pi}{2K'} (2hK \pm u) = \frac{q'^{-h} e^{\pm \frac{\pi u}{2K'}} - q'^h e^{\mp \frac{\pi u}{2K'}}}{q'^{-h} e^{\pm \frac{\pi u}{2K'}} + q'^h e^{\mp \frac{\pi u}{2K'}}};$$

$$\operatorname{sn} \frac{\pi}{2K'} (2hK + u) \operatorname{sn} \frac{\pi}{2K'} (2hK - u)$$

$$= \frac{q'^{-h} e^{\frac{\pi u}{2K'}} - q'^h e^{-\frac{\pi u}{2K'}}}{q'^{-h} e^{\frac{\pi u}{2K'}} + q'^h e^{-\frac{\pi u}{2K'}}} \cdot \frac{q'^{-h} e^{-\frac{\pi u}{2K'}} - q'^h e^{\frac{\pi u}{2K'}}}{q'^{-h} e^{-\frac{\pi u}{2K'}} + q'^h e^{\frac{\pi u}{2K'}}}$$

$$= \frac{q'^{-2h} + q'^{2h} - \left(e^{\frac{\pi u}{K'}} + e^{-\frac{\pi u}{K'}} \right)}{q'^{-2h} + q'^{2h} + \left(e^{\frac{\pi u}{K'}} + e^{-\frac{\pi u}{K'}} \right)}.$$

From plane trigonometry we have the equations

$$\frac{e^x - e^{-x}}{e^x + e^{-x}} = -i \tan ix, \quad e^x + e^{-x} = 2 \cos ix;$$

where $i = \sqrt{-1}$: which gives

$$\operatorname{sn} \frac{\pi u}{2K'} = -i \tan \frac{\pi i u}{2K'}; \quad (\text{Mod. } 1;)$$

$$\operatorname{sn} \frac{\pi}{2K'} (2hK + u) \operatorname{sn} \frac{\pi}{2K'} (2hK - u)$$

$$= \frac{q'^{-2h} + q'^{2h} - 2 \cos \frac{\pi i u}{K'}}{q'^{-2h} + q'^{2h} + 2 \cos \frac{\pi i u}{K'}}$$

$$= \frac{1 - 2q'^{2h} \cos \frac{\pi i u}{K'} + q'^{4h}}{1 + 2q'^{2h} \cos \frac{\pi i u}{K'} + q'^{4h}}.$$

From eq. (10), Chap. III, we have

$$\operatorname{sn} (u, k) = -i \operatorname{tn} (iu, k').$$

Substituting these values in eq. (6), we have

$$\operatorname{tn} (iu, k') = A' \tan \frac{\pi i u}{2K'} \left[\Pi \right] \frac{1 - 2q'^{2h} \cos \frac{\pi i u}{K'} + q'^{4h}}{1 + 2q'^{2h} \cos \frac{\pi i u}{K'} + q'^{4h}}.$$

Now in place of the series of moduli k', k'_0 and the corresponding complete integral K' , we are at liberty to substitute the parallel series of moduli k, k_0 and the corresponding complete integral K ; calling the new integral u , we have

$$(8) \quad \operatorname{tn} (u, k) = A \tan \frac{\pi u}{2K} \Pi \frac{1 - 2q^{2h} \cos \frac{\pi u}{K} + q^{4h}}{1 + 2q^{2h} \cos \frac{\pi u}{K} + q^{4h}}$$

$$\begin{aligned}
&= A \tan \frac{\pi u}{2K} \frac{1 - 2q^2 \cos \frac{\pi u}{K} + q^4}{1 + 2q^2 \cos \frac{\pi u}{K} + q^4} \\
&\quad \cdot \frac{1 - 2q^4 \cos \frac{\pi u}{K} + q^8}{1 + 2q^4 \cos \frac{\pi u}{K} + q^8} \\
&\quad \cdot \frac{1 - 2q^6 \cos \frac{\pi u}{K} + q^{12}}{1 + 2q^6 \cos \frac{\pi u}{K} + q^{12}} \dots,
\end{aligned}$$

where

$$(9) \quad A = (1 + k_0)(1 + k_{00})^2(1 + k_{02})^4(1 + k_{04})^8 \dots$$

Now in equation (6) put $u + K$ for u , and we have, since [eq. (II), Chap. III] $\operatorname{sn}(u + K) = \frac{\operatorname{cn} u}{\operatorname{dn} u}$,

$$\begin{aligned}
\frac{\operatorname{cn} u}{\operatorname{dn} u} &= A' \operatorname{sn} \frac{\pi(u + K)}{2K'} \left[II \right] \operatorname{sn} \frac{\pi}{2K'} [(2h + 1)K + u] \\
&\quad \cdot \operatorname{sn} \frac{\pi}{2K'} [(2h - 1)K - u]. \quad (\text{Mod. } 1.)
\end{aligned}$$

Now from $2h - 1$ and $2h + 1$ we have the following series of numbers respectively:

$$2h - 1: \quad 1, \quad 3, \quad 5, \quad 7, \quad 9, \quad \text{etc.}$$

$$2h + 1: \quad 3, \quad 5, \quad 7, \quad 9, \quad \text{etc.}$$

It will be observed that the factor outside of the sign $[II]$, viz., $\operatorname{sn} \operatorname{am} \frac{\pi(u + K)}{2K'}$, would, if placed under the sign $[II]$, supply

the missing first term of the second series. Hence, placing this factor within the sign, we have

$$(10) \quad \frac{\text{cn } u}{\text{dn } u} = A' [\Pi] \text{sn } \frac{\pi}{2K'} [(2h-1)K+u] \\ \cdot \text{sn } \frac{\pi}{2K'} [(2h-1)K-u]. \quad (\text{Mod. } 1.)$$

Comparing this with equation (7), we see that the factors herein differ from those in equation (7) only in having $2h-1$ in place of $2h$; hence we have

$$\text{sn } \frac{\pi}{2K'} [(2h-1)K+u] \text{sn } \frac{\pi}{2K'} [(2h-1)K-u] \\ = \frac{1 - 2q'^{2h-1} \cos \frac{\pi iu}{K'} + q'^{4h-2}}{1 + 2q'^{2h-1} \cos \frac{\pi iu}{K'} + q'^{4h-2}}. \quad (\text{Mod. } 1.)$$

From eqs. (10), Chap. III, we have

$$\frac{\text{cn } (u, k)}{\text{dn } (u, k)} = \frac{1}{\text{dn } (iu, k')};$$

whence eq. (10) becomes

$$(11) \quad \frac{1}{\text{dn } (iu, k')} = A' [\Pi] \frac{1 - 2q'^{2h-1} \cos \frac{\pi iu}{K'} + q'^{4h-2}}{1 + 2q'^{2h-1} \cos \frac{\pi iu}{K'} + q'^{4h-2}};$$

and when in place of iu, k', K', q', A' , we substitute u, k, K, q and A , and invert the equation, we have

$$(12) \quad \text{dn } (u, k) = \frac{1}{A} [\Pi] \frac{1 + 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}}{1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}}.$$

Bearing in mind the remarkable property (Chap. III, p. 29) that the functions $\text{sn } u$ and $\text{dn } u$ approach infinity for the same value of u , we see that both these functions, except as to the factor independent of u , must have the same denominator. Furthermore, since $\text{sn } u$ and $\text{tn } u$ disappear for the same value of u , they must, except for the independent factor, have the same numerator. Hence, indicating by B a new quantity, dependent upon k but independent of u , we have

$$(13) \quad \text{sn } (u, k) = B \sin \frac{\pi u}{2K} \left[\Pi \right] \frac{1 - 2q^{2h} \cos \frac{\pi u}{K} + q^{4h}}{1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}} ;$$

and since

$$\text{cn } u = \frac{\text{sn } u}{\text{tn } u},$$

we also have, from eqs. (8) and (13),

$$(14) \quad \text{cn } (u, k) = \frac{B}{A} \cos \frac{\pi u}{2K} \left[\Pi \right] \frac{1 + 2q^{2h} \cos \frac{\pi u}{K} + q^{4h}}{1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}}.$$

Collecting these results, we have the following equations :

$$(15) \quad \text{sn } (u, k) = B \sin \frac{\pi u}{2K} \left[\Pi \right] \frac{1 - 2q^{2h} \cos \frac{\pi u}{K} + q^{4h}}{1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}},$$

$$(16) \quad \text{cn } (u, k) = \frac{B}{A} \cos \frac{\pi u}{2K} \left[\Pi \right] \frac{1 + 2q^{2h} \cos \frac{\pi u}{K} + q^{4h}}{1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}},$$

$$(17) \quad \operatorname{dn}(u, k) = \frac{1}{A} \left[\Pi \right] \frac{1 + 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}}{1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}}.$$

To ascertain the values of A and B , we proceed as follows:

In eq. (17) we make $u = 0$, whence, by eq. (13), Chap. II, we have

$$1 = \frac{1}{A} \left[\Pi \right] \left(\frac{1 + q^{2h-1}}{1 - q^{2h-1}} \right)^2;$$

whence

$$(18) \quad \frac{1}{A} = \left[\Pi \right] \left(\frac{1 - q^{2h-1}}{1 + q^{2h-1}} \right)^2.$$

In equation (17), making $u = K$, we get, by equation (1), Chap. III,

$$k' = \frac{1}{A} \left[\Pi \right] \left(\frac{1 - q^{2h-1}}{1 + q^{2h-1}} \right)^2 = \frac{1}{A^2};$$

$$(19) \quad \therefore \frac{1}{A} = \sqrt{k'}.$$

whence

$$\frac{B}{A} = B \sqrt{k'}.$$

To calculate B , put $e^{\frac{i\pi u}{2K}} = v$; if we change $\frac{\pi u}{2K}$ into

$\frac{\pi u}{2K} + \frac{i\pi K'}{2K}$, v will change into $v\sqrt{q}$, and $\text{sn } u$ will become, by eq. (14), Chap. III,

$$\text{sn}(u + iK') = \frac{1}{k \text{sn } u}.$$

Now, replacing $\sin \frac{\pi u}{2K}$ and $\cos \frac{\pi u}{K}$ by their exponential values, and observing that

$$1 - 2q^n \cos \frac{\pi u}{K} + q^{2n} = (1 - q^n v^3)(1 - q^n v^{-2}),$$

we have

$$\text{sn } u = \frac{B}{2} \cdot \frac{v - v^{-1}}{\sqrt{1 - 1}} \cdot \frac{[II](1 - q^{2h} v^3)(1 - q^{2h} v^{-2})}{[II](1 - q^{2h-1} v^3)(1 - q^{2h-1} v^{-2})}.$$

Changing u into $u + iK'$, and consequently v into $v\sqrt{q}$, we have

$$\frac{1}{k \text{sn } u} = \frac{B}{2} \cdot \frac{v\sqrt{q} - v^{-1}\sqrt{q^{-1}}}{\sqrt{1 - 1}} \cdot \frac{[II](1 - q^{2h+1} v^3)(1 - q^{2h-1} v^{-2})}{[II](1 - q^{2h} v^3)(1 - q^{2h-2} v^{-2})}.$$

Multiplying these equations together, member by member, and observing that

$$v\sqrt{q} - v^{-1}\sqrt{q^{-1}} = \frac{1 - qv^3}{-v\sqrt{q}},$$

$$v - v^{-1} = v(1 - v^{-2}),$$

we get

$$\frac{1}{k} = \frac{B^2}{4} \cdot \frac{1 - qv^3}{v\sqrt{q}} \cdot v(1 - v^{-2}) \cdot \frac{[II](1 - q^{2h+1} v^3)(1 - q^{2h} v^{-2})}{[II](1 - q^{2h-1} v^3)(1 - q^{2h-2} v^{-2})}$$

$$\begin{aligned}
&= \frac{B^2}{4\sqrt{q}} (1 - q\nu^2) (1 - \nu^{-2}) \frac{(1 - q^3\nu^2)(1 - q^5\nu^2) \dots}{(1 - q\nu^2)(1 - q^3\nu^2) \dots} \\
&\quad \cdot \frac{(1 - q^2\nu^{-2})(1 - q^4\nu^{-2}) \dots}{(1 - \nu^{-2})(1 - q^2\nu^{-2}) \dots} \\
&= \frac{B^2}{4} \cdot \frac{1}{\sqrt{q}}.
\end{aligned}$$

$$\therefore B = \frac{2\sqrt[4]{q}}{\sqrt{k}};$$

whence

$$\frac{B}{A} = 2\sqrt[4]{q}\sqrt{\frac{k'}{k}}.$$

Substituting these values in eqs. (15), (16), and (17), we have

$$(20) \quad \operatorname{sn}(u, k) = \frac{2\sqrt[4]{q}}{\sqrt{k}} \sin \frac{\pi u}{2K} \left[\Pi \right] \frac{1 - 2q^{2h} \cos \frac{\pi u}{K} + q^{4h}}{1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}};$$

$$(21) \quad \operatorname{cn}(u, k) = \frac{2\sqrt{k'}\sqrt[4]{q}}{\sqrt{k}} \cos \frac{\pi u}{2K} \left[\Pi \right] \frac{1 + 2q^{2h} \cos \frac{\pi u}{K} + q^{4h}}{1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}};$$

$$(22) \quad \operatorname{dn}(u, k) = \sqrt{k'} \left[\Pi \right] \frac{1 + 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}}{1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}}.$$

CHAPTER VIII.

THE Θ FUNCTION.

WE will indicate the denominator in eq. (20), Chap. VII, by $\phi(u)$, thus:

$$(1) \quad \phi(u) = [\Pi](1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}).$$

We will now develop this into a series consisting of the cosines of the multiples of $\frac{\pi u}{K}$. Put $\frac{\pi u}{2K} = x$, whence

$$2 \cos \frac{\pi u}{K} = (e^{2ix} + e^{-2ix});$$

but

$$1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2} = (1 - q^{2h-1} e^{2ix})(1 - q^{2h-1} e^{-2ix}),$$

and therefore

$$(2) \quad \begin{aligned} \phi(u) &= (1 - qe^{2ix})(1 - q^3e^{2ix})(1 - q^5e^{2ix}) \dots \\ &\quad (1 - qe^{-2ix})(1 - q^3e^{-2ix})(1 - q^5e^{-2ix}) \dots \end{aligned}$$

Putting now $u + 2iK'$ instead of u , we have

$$x_1 = \frac{\pi(u + 2iK')}{2K} = x + \frac{\pi iK'}{K},$$

$$2ix_1 = 2ix - \frac{2\pi K'}{K};$$

and

$$e^{2ix_1} = q^2 e^{2ix},$$

$$e^{-2ix_1} = \frac{1}{q^2} e^{-2ix}.$$

From these we have

$$\phi(u + 2iK') = -\frac{1}{q} e^{-2ix} (1 - qe^{2ix})(1 - q^3 e^{2ix}) \dots$$

$$(1 - qe^{-2ix})(1 - q^3 e^{-2ix}) \dots;$$

whence

$$\phi(u + 2iK') = -\frac{1}{q} e^{-2ix} \phi(u),$$

or

$$(3) \quad \phi(u + 2iK') = -q^{-1} e^{-\frac{\pi i u}{K}} \phi(u).$$

Now put

$$(4) \quad \phi(u) = A + B \cos \frac{\pi u}{K} + C \cos \frac{2\pi u}{K} + D \cos \frac{3\pi u}{K} + \text{etc.}$$

Since

$$\cos \frac{\pi u}{K} = \frac{1}{2}(e^{2ix} + e^{-2ix}),$$

this becomes

$$(5) \quad \phi(u) = A + \frac{1}{2} B e^{2ix} + \frac{1}{2} C e^{4ix} + \frac{1}{2} D e^{6ix} + \dots$$

$$+ \frac{1}{2} B e^{-2ix} + \frac{1}{2} C e^{-4ix} + \frac{1}{2} D e^{-6ix} + \dots;$$

whence

$$(6) \quad -\frac{1}{q} e^{-2ix} \phi(u) = -\frac{A}{q} e^{-2ix} - \frac{B}{2q} - \frac{C}{2q} e^{2ix} - \frac{D}{2q} e^{4ix} - \dots$$

$$- \frac{B}{2q} e^{-4ix} - \frac{C}{2q} e^{-6ix} - \frac{D}{2q} e^{-8ix} - \dots$$

Now in equation (5) put $u + 2iK'$ in place of u , remembering that e^{2ix} and e^{-2ix} are thereby changed respectively into $q^2 e^{2ix}$ and $q^{-2} e^{-2ix}$, and we have

$$(7) \quad \phi(u + 2iK') = A + \frac{Bq^2}{2} e^{2ix} + \frac{Cq^4}{2} e^{4ix} + \frac{Dq^6}{2} e^{6ix} + \dots \\ + \frac{B}{2q^2} e^{-2ix} + \frac{C}{2q^4} e^{-4ix} + \dots$$

Since equations (6) and (7) are equal, we have

$$-\frac{B}{2q} = A, \quad B = -2qA;$$

$$-\frac{C}{2q} = \frac{Bq^2}{2}, \quad C = +2q^3 A;$$

$$-\frac{D}{2q} = \frac{Cq^2}{2}, \quad D = -2q^5 A;$$

$$\dots \dots \dots$$

whence

$$(8) \quad \left\{ \begin{aligned} & [II](1 - 2q^{2k-1} \cos \frac{\pi u}{K} + q^{4k-2}) \\ & = A(1 - 2q \cos \frac{\pi u}{K} + 2q^4 \cos \frac{2\pi u}{K} - 2q^9 \cos \frac{3\pi u}{K} \\ & \quad + 2q^{16} \cos \frac{4\pi u}{K} - \dots). \end{aligned} \right.$$

The series in the second member has been designated by Jacobi and subsequent writers by $\Theta(u)$, thus:

$$(9) \quad \Theta(u) = 1 - 2q \cos \frac{\pi u}{K} + 2q^4 \cos \frac{2\pi u}{K} - \dots$$

CHAPTER IX.

THE Θ AND H FUNCTIONS.

IN equation (20), Chap. VII, viz.,

$$\operatorname{sn}(u, k) = \frac{2\sqrt[4]{q}}{\sqrt{k}} \sin \frac{\pi u}{2K} \left[\Pi \right] \frac{1 - 2q^{2h} \cos \frac{\pi u}{K} + q^{4h}}{1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}},$$

the numerator and the denominator have been considered separately by Jacobi, who gave them a special notation and developed from them a theory second only in importance to the elliptic functions themselves.

Put [see equation (8), Chap. VIII]

$$(1) \quad \Theta(u) = \frac{1}{A} \left[\Pi \right] (1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}).$$

$$(2) \quad H(u) = 2 \frac{1}{A} \sqrt[4]{q} \sin \frac{\pi u}{2K} \left[\Pi \right] (1 - 2q^{2h} \cos \frac{\pi u}{K} + q^{4h});$$

A being a constant whose value is to be determined later. From these we have

$$(3) \quad \operatorname{sn}(u, k) = \frac{1}{\sqrt{k}} \cdot \frac{H(u)}{\Theta(u)}.$$

The functions $\operatorname{sn} u$ and $\operatorname{cn} u$ can also be expressed in terms of the new functions; thus we have

$$(4) \quad \operatorname{cn}(u, k) = \sqrt{\frac{k'}{k}} \cdot 2\sqrt{q} \cos \frac{\pi u}{2K} \left[\Pi \right] \frac{1 + 2q^{2h} \cos \frac{\pi u}{K} + q^{4h}}{1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}};$$

or, since $\sin x = \cos \left(x + \frac{\pi}{2} \right)$ and $\cos x = -\cos \left(x + \frac{\pi}{2} \right)$,

and putting $u = \frac{2Kx}{\pi}$,

$$\begin{aligned} \operatorname{cn} \left(\frac{2Kx}{\pi}, k \right) &= \sqrt{\frac{k'}{k}} \frac{H \left[\frac{2K}{\pi} \left(x + \frac{\pi}{2} \right) \right]}{\Theta \left(\frac{2Kx}{\pi} \right)} \\ &= \sqrt{\frac{k'}{k}} \frac{H \left[\frac{2Kx}{\pi} + K \right]}{\Theta \left(\frac{2Kx}{\pi} \right)}. \end{aligned}$$

Replacing $\frac{2Kx}{\pi}$ by its value, u , we have

$$(5) \quad \operatorname{cn}(u, k) = \sqrt{\frac{k'}{k}} \frac{H(u + K)}{\Theta(u)}.$$

Furthermore,

$$(6) \quad \operatorname{dn}(u, k) = \sqrt{k'} \left[\Pi \right] \frac{1 + 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}}{1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}}.$$

gives in the same manner

$$\operatorname{dn} \frac{2Kx}{\pi} = \sqrt{k'} \frac{\Theta \left[\frac{2K}{\pi} \left(x + \frac{\pi}{2} \right) \right]}{\Theta \left(\frac{2Kx}{\pi} \right)},$$

or

$$(7) \quad \operatorname{dn} (u, k) = \sqrt{k'} \frac{\Theta(u + K)}{\Theta(u)}.$$

If we put

$$(8) \quad H(u + K) = H_1(u),$$

$$(9) \quad \Theta(u + K) = \Theta_1(u),$$

the three elliptic functions can be expressed by the following formulas:

$$(10) \quad \operatorname{sn} (u, k) = \frac{1}{\sqrt{k}} \cdot \frac{H(u)}{\Theta(u)};$$

$$(11) \quad \operatorname{cn} (u, k) = \sqrt{\frac{k'}{k}} \cdot \frac{H_1(u)}{\Theta(u)};$$

$$(12) \quad \operatorname{dn} (u, k) = \sqrt{k'} \frac{\Theta_1(u)}{\Theta(u)}.$$

These functions Θ and H can be expressed in terms of each other. By definition,

$$H(u) = 2C \sqrt{q} \sin \frac{\pi u}{2K} \left[\Pi \right] \left(1 - 2q^{2h} \cos \frac{\pi u}{K} + q^{4h} \right);$$

but

$$1 - 2q^h \cos \frac{\pi u}{K} + q^{2h} = \left(1 - q^h e^{\frac{\pi u \sqrt{-1}}{K}}\right) \left(1 - q^h e^{-\frac{\pi u \sqrt{-1}}{K}}\right),$$

$$\sin \frac{\pi u}{2K} = \frac{e^{\frac{\pi i u}{2K}} - e^{-\frac{\pi i u}{2K}}}{2 \sqrt{-1}}$$

$$= e^{\frac{-\pi i u}{2K}} \frac{1 - e^{\frac{\pi i u}{K}}}{2} \sqrt{-1},$$

and consequently

$$(13) \quad H(u) = C \sqrt[4]{q} e^{-\frac{\pi i u}{2K}} \sqrt{-1} \left(1 - e^{\frac{\pi i u}{K}}\right) \left(1 - q^2 e^{-\frac{\pi i u}{K}}\right) \left(1 - q^2 e^{\frac{\pi i u}{K}}\right) \dots$$

Now, changing u into $u + iK'$, and remembering that $e^{\frac{-\pi K'}{K}} = q$, we have

$$(14) \quad H(u + iK')$$

$$= C q^{-\frac{1}{4}} e^{-\frac{\pi i u}{2K}} \sqrt{-1} \left(1 - q e^{\frac{\pi i u}{K}}\right) \left(1 - q e^{-\frac{\pi i u}{K}}\right) \left(1 - q^3 e^{\frac{\pi i u}{K}}\right) \left(1 - q^3 e^{-\frac{\pi i u}{K}}\right) \dots;$$

and reuniting the factors two by two, this becomes

$$(15) \quad H(u + iK')$$

$$= C \sqrt{-1} q^{-\frac{1}{4}} e^{-\frac{\pi i u}{2K}} \left(1 - 2q \cos \frac{\pi u}{K} + q^2\right) \left(1 - 2q^3 \cos \frac{\pi u}{K} + q^6\right) \dots;$$

and finally, according to equation (1),

$$(16) \quad H(u + iK') = \sqrt{-1} q^{-\frac{1}{4}} e^{-\frac{\pi i u}{2K}} \Theta(u).$$

In the same manner, we can get

$$(17) \quad \Theta(u + iK') = \sqrt{-1} q^{-\frac{1}{2}} e^{-\frac{\pi i u}{2K}} H(u).$$

Substituting $u + 2K$ for u in equations (1) and (2), we get

$$(18) \quad \Theta(u + 2K) = \Theta(u),$$

$$(19) \quad H(u + 2K) = -H(u),$$

$$\text{since } \cos \frac{\pi}{K}(u + 2K) = \cos \frac{\pi u}{K} \text{ and } \sin \frac{\pi}{2K}(u + 2K) = -\sin \frac{\pi u}{2K}.$$

The comparison of these four equations with equations (10), (11), and (12) shows the periodicity of the elliptic functions. For example, comparing eqs. (10) and (16) and (17), we see that changing u into $u + iK'$ simply multiplies the numerator and denominator of the second member of eq. (10) by the same number, and does not change their ratio.

The addition of $2K$ changes the sign of the function, but not its value.

We will define Θ_1 and H_1 as follows:

$$(20) \quad \Theta_1(x) = \Theta(x + K);$$

$$(21) \quad H_1(x) = H(x + K).$$

Hence we get, from equation (17),

$$\begin{aligned} \Theta_1(x + iK') &= \Theta(x + iK' + K) = \Theta(x + K + iK') \\ &= iH(x + K)e^{-\frac{i\pi}{4K}(2x + 2K + iK')} \\ &= iH_1(x)e^{-\frac{i\pi}{4K}(2x + iK')}(-\sqrt{-1}), \end{aligned}$$

$$\text{since } e^{-\frac{i\pi}{2}} = \cos \frac{\pi}{2} - \sqrt{-1} \sin \frac{\pi}{2} = -\sqrt{-1};$$

whence

$$(22) \quad \Theta_1(x + iK') = H_1(x) e^{-\frac{i\pi}{4K}(2x + iK')}.$$

In a similar manner we get

$$(22)^* \quad H_1(x + iK') = \Theta_1(x) e^{-\frac{i\pi}{4K}(2x + iK')}.$$

In eq. (9), Chap. VIII, put $u = \frac{2Kz}{\pi}$, and we get

$$(23) \quad \Theta\left(\frac{2Kz}{\pi}\right) = 1 - 2q \cos 2z + 2q^4 \cos 4z - \dots$$

Now, in this equation, changing z into $z + \frac{\pi}{2}$, and observing eq. (20), we get

$$(24) \quad \Theta_1\left(\frac{2Kz}{\pi}\right) = 1 + 2q \cos 2z + 2q^4 \cos 4z + \dots$$

Applying eq. (22) to this, we have

$$\begin{aligned} H_1\left(\frac{2Kz}{\pi}\right) &= \Theta_1\left(\frac{2K}{\pi}\left(z + \frac{\pi iK'}{2K}\right)\right) e^{\frac{\pi i}{4K}\left(\frac{4Kz}{\pi} + iK'\right)} \\ &= \Theta_1\left(\frac{2K}{\pi}\left(z + \frac{\pi iK'}{2K}\right)\right) e^{iz} q^{\frac{1}{4}} \\ &= e^{iz} q^{\frac{1}{4}} \left[1 + 2q \cos 2\left(z + \frac{\pi iK'}{2K}\right) + 2q^4 \cos 4\left(z + \frac{\pi iK'}{2K}\right) + \dots \right] \\ &= e^{iz} q^{\frac{1}{4}} \left[1 + q \left(e^{2i\left(z + \frac{\pi iK'}{2K}\right)} + e^{-2i\left(z + \frac{\pi iK'}{2K}\right)} \right) \right. \\ &\quad \left. + q^4 \left(e^{4i\left(z + \frac{\pi iK'}{2K}\right)} + e^{-4i\left(z + \frac{\pi iK'}{2K}\right)} \right) + \dots \right] \end{aligned}$$

$$\begin{aligned}
&= e^{iz} q^{\frac{1}{2}} [1 + q(qe^{2iz} + q^{-1}e^{-2iz}) + q^4(q^2e^{4iz} + q^{-2}e^{-4iz}) + \dots] \\
&= e^{iz} q^{\frac{1}{2}} [1 + q^2e^{2iz} + q^0e^{4iz} + \dots \\
&\quad + e^{-2iz} + q^3e^{-4iz} + \dots] \\
&= q^{\frac{1}{2}} [e^{iz} + q^2e^{3iz} + q^0e^{5iz} + \dots \\
&\quad + e^{-iz} + q^3e^{-3iz} + q^0e^{-5iz} + \dots] \\
&= 2q^{\frac{1}{2}} [\cos z + q^2 \cos 3z + q^0 \cos 5z + \dots];
\end{aligned}$$

whence

$$(25) \quad H_1\left(\frac{2Kz}{\pi}\right) = 2\sqrt[4]{q} \cos z + 2\sqrt[4]{q^9} \cos 3z + 2\sqrt[4]{q^{25}} \cos 5z + \dots$$

In this equation, changing z into $z - \frac{\pi}{2}$, and applying eq. (21), we get

$$(26) \quad H\left(\frac{2Kz}{\pi}\right) = 2\sqrt[4]{q} \sin z - 2\sqrt[4]{q^9} \sin 3z + 2\sqrt[4]{q^{25}} \sin 5z - \dots,$$

since

$$H_1\left(\frac{2Kz}{\pi}\right) = H\left(\frac{2Kz}{\pi} + K\right).$$

We will now determine the constant A of eq. (8), Chap. VIII, and eqs. (1) and (2) of this chapter. Denote A by $f(q)$, and we have

$$(26)^* \quad [II](1 - 2q^{2h-1} \cos \frac{\pi u}{K} + q^{4h-2}) = f(q)\Theta(u).$$

Substituting herein $u = 0$ and $u = \frac{K}{2}$, we have

$$[II](1 - q^{2h-1})^2 = f(q)\Theta(0);$$

$$[II](1 + q^{4h-2}) = f(q)\Theta\left(\frac{K}{2}\right).$$

From eq. (9), Chap. VIII, we get

$$(27) \quad \Theta(0) = 1 - 2q + 2q^4 - 2q^9 + 2q^{16} - \dots;$$

$$(28) \quad \Theta\left(\frac{K}{2}\right) = 1 - 2q^4 + 2q^{16} - 2q^{36} + 2q^{64} - \dots;$$

from which we see that $\Theta(0)$ is changed into $\Theta\left(\frac{K}{2}\right)$ when we put q^4 in place of q .

Whence

$$[\Pi](1 - q^{8h-4})^2 = f(q^4)\Theta\left(\frac{K}{2}\right);$$

and therefore

$$(29) \quad \frac{f(q)}{f(q^4)} = [\Pi] \frac{1 + q^{4h-2}}{(1 - q^{8h-4})^2} \\ = [\Pi] \frac{1}{(1 - q^{8h-4})(1 - q^{4h-2})}.$$

Now, the expressions $4h - 2$, $8h - 4$, and $8h$ give the following series of numbers:

$$\begin{array}{cccccccc} 4h - 2, & & 2, & 6, & 10, & 14, & 18, & 22, & 26, & 30, & 34; \\ 8h - 4, & & & 4, & & 12, & & 20, & & 28, & & 36; \\ 8h, & & & & & 8, & & 16, & & 24, & & 32. \end{array}$$

Hence, the three expressions taken together contain all the even numbers, and

$$[\Pi](1 - q^{8h-4})(1 - q^{4h-2})(1 - q^{8h}) = [\Pi](1 - q^{2h}).$$

Therefore, multiplying eq. (29) by

$$[\Pi] \frac{1 - q^{8h}}{1 - q^{2h}},$$

we have

$$\frac{f(q)}{f(q^4)} = [\Pi] \frac{1 - q^{8h}}{1 - q^{2h}}.$$

Now in this equation, by successive substitutions of q^4 for q , we get

$$\frac{f(q^4)}{f(q^{16})} = \left[\Pi \right] \frac{1 - q^{32h}}{1 - q^{8h}};$$

$$\frac{f(q^{16})}{f(q^{64})} = \left[\Pi \right] \frac{1 - q^{128h}}{1 - q^{32h}};$$

$$\frac{f(q^{64})}{f(q^{256})} = \left[\Pi \right] \frac{1 - q^{512h}}{1 - q^{128h}};$$

.

Now q being less than 1, q^n tends towards the limit 0 as n increases, and consequently $1 - q^n$ tends towards the limit 1. Also, from eq. (8), Chap. VIII, we see that $f(0) = 1$. Hence, multiplying the above equations together member by member, we have

$$(30) \quad f(q) = \left[\Pi \right] \frac{1}{1 - q^{2h}},$$

or

$$(31) \quad A = \frac{1}{(1 - q^2)(1 - q^4)(1 - q^6) \dots}.$$

Substituting this value in equation (8), Chap. VIII, we have, after making $u = 0$,

$$\begin{aligned} (1 - q)^2(1 - q^2)^2(1 - q^4)^2 \dots &= \frac{1 - 2q + 2q^4 - 2q^9 + \dots}{(1 - q^2)(1 - q^4)(1 - q^6) \dots} \\ &= \frac{\Theta(0)}{(1 - q^2)(1 - q^4)(1 - q^6) \dots}. \end{aligned}$$

(See equation (9), Chap. VIII.)

Transposing one of the series of products from the left-hand member, we get

$$(1 - q)(1 - q^3) \dots = \frac{\Theta(0)}{(1 - q)(1 - q^2)(1 - q^3)(1 - q^4) \dots}.$$

Introducing on both sides of the equation the factors $1 - q^2$, $1 - q^4$, $1 - q^6$, etc., we get

$$\begin{aligned} & (1 - q)(1 - q^2)(1 - q^3)(1 - q^4) \dots \\ &= \Theta(0) \frac{1 - q^2}{1 - q} \cdot \frac{1 - q^4}{1 - q^2} \cdot \frac{1 - q^6}{1 - q^3} \cdot \frac{1 - q^8}{1 - q^4} \dots \\ &= \Theta(0)(1 + q)(1 + q^2)(1 + q^3) \dots; \end{aligned}$$

whence

$$(32) \quad \Theta(0) = \frac{(1 - q)(1 - q^2)(1 - q^3)}{(1 + q)(1 + q^2)(1 + q^3)}.$$

Resuming equation (20), Chap. VII, and dividing both members of the equation by u , we have

$$\frac{\operatorname{sn} u}{u} = \frac{2\sqrt[4]{q}}{\sqrt{k}} \cdot \frac{\sin \frac{\pi u}{2K}}{u} \left[II \right] \frac{1 - 2q^{2k} \cos \frac{\pi u}{K} + q^{4k}}{1 - 2q^{2k-1} \cos \frac{\pi u}{K} + q^{4k-2}}.$$

This, for $u = 0$, since the limiting value of $\frac{\operatorname{sn} u}{u}$ for $u = 0$ is 1,

and of $\frac{\sin \frac{\pi u}{2K}}{u}$ for $u = 0$ is $\frac{\pi}{2K}$, becomes

$$1 = \frac{\sqrt[4]{q}}{\sqrt{k}} \cdot \frac{\pi}{K} \cdot \frac{(1 - q^2)^2(1 - q^4)^2(1 - q^6)^2 \dots}{(1 - q)^2(1 - q^3)^2(1 - q^5)^2 \dots},$$

or

$$(33) \quad \frac{\sqrt{k}K}{\pi\sqrt[4]{q}} = \left[\frac{(1 - q^2)(1 - q^4)(1 - q^6) \dots}{(1 - q)(1 - q^3)(1 - q^5) \dots} \right]^2.$$

Further, from equation (21), Chap. VII, for $u = 0$, we have

$$(34) \quad \frac{\sqrt{k}}{2\sqrt{k'}\sqrt[4]{q}} = \left[\frac{(1 + q^2)(1 + q^4)(1 + q^6) \dots}{(1 - q)(1 - q^3)(1 - q^5) \dots} \right]^2.$$

The quotient of these two equations gives

$$(35) \quad \frac{2\sqrt{k'}K}{\pi} = \left[\frac{(1-q^2)(1-q^4)(1-q^6)\dots}{(1+q^2)(1+q^4)(1+q^6)\dots} \right]^2;$$

or, substituting the value of $\sqrt{k'}$ from eqs. (18) and (19), Chap. VII,

$$(36) \quad \frac{2k'K}{\pi} = \left[\frac{(1-q)(1-q^2)(1-q^3)\dots}{(1+q)(1+q^2)(1+q^3)\dots} \right]^2.$$

Comparing this with equation (32), we easily get

$$(37) \quad \Theta(0) = \sqrt{\frac{2k'K}{\pi}}.$$

From equation (9), Chap. VIII, making $u = K$, we get

$$(38) \quad \Theta(K) = 1 + 2q + 2q^4 + 2q^9 + 2q^{16} + \dots$$

Making $z = 0$ in equation (24), Chap. IX, we have

$$(39) \quad \Theta_1(0) = 1 + 2q + 2q^4 + 2q^9 + \dots$$

This might also have been derived from eq. (38) by observing that

$$\Theta(0+K) = \Theta_1(0) = \Theta(K).$$

Knowing $\Theta(0)$, it is easy to deduce $\Theta(K)$ and $H(K)$

From equation (7) we have

$$\operatorname{dn} u = \sqrt{k'} \frac{\Theta(u+K)}{\Theta(u)}.$$

Making $u = 0$, we have, since $\operatorname{dn}(0) = 1$,

$$(40) \quad \Theta(K) = \frac{\Theta(0)}{\sqrt{k'}}.$$

From equation (5) we get, in the same manner,

$$(41) \quad H(K) = \sqrt{\frac{k'}{k}} \mathcal{E}(0).$$

From eq. (12), Chap. IX, we have

$$(41)^* \quad \operatorname{dn} u = \sqrt{1 - k^2 \sin^2 \phi} = \sqrt{k'} \frac{\mathcal{E}_1(u)}{\mathcal{E}(u)};$$

and putting $x = \frac{\pi u}{2K}$, we have

$$(42) \quad \frac{\operatorname{dn} u}{\sqrt{k'}} = \frac{1 + 2q \cos 2x + 2q^4 \cos 4x + 2q^9 \cos 6x + \dots}{1 - 2q \cos 2x + 2q^4 \cos 4x - 2q^9 \cos 6x + \dots}.$$

Putting

$$(42)^* \quad \frac{\operatorname{dn} u}{\sqrt{k'}} = \cot \gamma,$$

we have

$$\frac{\cot \gamma - 1}{\cot \gamma + 1} = \tan(45^\circ - \gamma) = 2q \frac{\cos 2x + q^4(4 \cos^2 2x - 3 \cos 2x) + \dots}{1 + q^4(4 \cos^2 2x - 2)};$$

whence

$$(43) \quad \cos 2x = \frac{\tan(45^\circ - \gamma)[1 + q^4(4 \cos^2 2x - 2)]}{2q - q^4(4 \cos^2 2x - 3 \cos 2x) - \dots},$$

and approximately,

$$(44) \quad \cos 2x = \frac{\tan(45^\circ - \gamma)}{2q}.$$

From equations (37) and (40), Chap. IX, we have

$$(45) \quad x = \frac{u}{\Theta^2(K)};$$

whence

$$(46) \quad u = x\Theta^2(K).$$

CHAPTER X.

ELLIPTIC INTEGRALS OF THE SECOND ORDER.

FROM Chap. I, equation (19), we have

$$E(k, \phi) = \int_0^\phi \sqrt{1 - k^2 \sin^2 \phi} \cdot d\phi = \int_0^\phi \Delta\phi \cdot d\phi.$$

From this we have

$$E(\phi) + E(\psi) = \int_0^\phi \Delta\phi \cdot d\phi + \int_0^\psi \Delta\phi \cdot d\phi.$$

Put

$$(1) \quad E\phi + E\psi = S.$$

Differentiating, we get

$$(2) \quad \Delta\phi \cdot d\phi + \Delta\psi \cdot d\psi = dS.$$

But we have, Chap. II, equation (2),

$$\frac{d\phi}{\Delta\phi} + \frac{d\psi}{\Delta\psi} = 0,$$

or

$$(3) \quad \Delta\psi \cdot d\phi + \Delta\phi \cdot d\psi = 0.$$

Adding equations (2) and (3), we get

$$(4) \quad (\Delta\phi + \Delta\psi)(d\phi + d\psi) = dS.$$

Substituting $\cos \mu$ from eq. (5), in eq. (5)*, Chap. II, we get

$$(5) \quad \begin{cases} \Delta\phi = \frac{\sin \phi \cos \psi \Delta\mu + \cos \phi \sin \psi}{\sin \mu}, \\ \Delta\psi = \frac{\sin \psi \cos \phi \Delta\mu + \cos \psi \sin \phi}{\sin \mu}; \end{cases}$$

whence

$$(6) \quad \Delta\phi \pm \Delta\psi = \frac{\Delta\mu \pm 1}{\sin \mu} \sin(\phi \pm \psi).$$

Substituting in equation (4), we have

$$(7) \quad \begin{aligned} dS &= \frac{\Delta\mu + 1}{\sin \mu} \sin(\phi + \psi) d(\phi + \psi) \\ &= -\frac{\Delta\mu + 1}{\sin \mu} d \cos(\phi + \psi). \end{aligned}$$

Integrating equation (7), we have

$$E\phi + E\psi = \frac{\Delta\mu + 1}{\sin \mu} [C - \cos(\phi + \psi)].$$

The constant of integration, C , is determined by making $\phi = 0$; in this case $\psi = \mu$, $E\phi = 0$, $E\psi = E\mu$, and $S = E\mu$; whence

$$E\mu = \frac{\Delta\mu + 1}{\sin \mu} (C - \cos \mu),$$

and by subtraction,

$$E\phi + E\psi - E\mu = \frac{\Delta\mu + 1}{\sin \mu} (\cos \mu - \cos \phi \cos \psi + \sin \phi \sin \psi).$$

But, Chap. II, eq. (5),

$$\cos \mu - \cos \phi \cos \psi = -\sin \phi \sin \psi \Delta\mu;$$

whence

$$E\phi + E\psi - E\mu = \frac{1 - k^2 \sin^2 \mu}{\sin \mu} \sin \phi \sin \psi$$

whence

$$(8) \quad E\phi + E\psi = E\mu + k^2 \sin \phi \sin \psi \sin \mu.$$

When $\phi = \psi$, we have

$$(9) \quad E\mu = 2E\phi - k^2 \sin^2 \phi \sin \mu.$$

But in that case

$$(10) \quad \cos \mu = \cos^2 \phi - \sin^2 \phi \Delta \mu;$$

whence

$$(11) \quad \sin \phi = \sqrt{\frac{1 - \cos \mu}{1 + \Delta \mu}}.$$

Let ϕ, ϕ_1, ϕ_2 , etc., be such values as will satisfy the equations

$$(12) \quad \begin{aligned} E\phi &= 2E\phi_1 - k^2 \sin^2 \phi_1 \sin \phi, \\ E\phi_1 &= 2E\phi_2 - k^2 \sin^2 \phi_2 \sin \phi_1, \\ &\text{etc.} \qquad \qquad \text{etc.} \end{aligned}$$

Assume an auxiliary angle γ , such that

$$(13) \quad \sin \gamma = k \sin \phi;$$

whence

$$\Delta \phi = \cos \gamma,$$

and Chap. IV, eq. (24),

$$(14) \quad \sin \phi_1 = \frac{\sin \frac{1}{2}\phi}{\cos \frac{1}{2}\gamma}.$$

Substituting in eq. (16) the numerical values derived from equations (15) and (17), we are enabled to determine the value of $E\phi$.

Landen's Transformation can also be applied to Elliptic Integrals of this class.

From eq. (11), Chap. IV, we get, by easy transformation,

$$(18) \quad \sin^2 2\phi = \sin^2 \phi_1 (1 + k_0 + 2k_0 \cos 2\phi).$$

From this we easily get

$$\begin{aligned} 2k_0 \cos 2\phi \sin^2 \phi_1 &= \sin^2 2\phi - \sin^2 \phi_1 - k_0^2 \sin^2 \phi_1 \\ &= 1 - \cos^2 2\phi - \sin^2 \phi_1 - k_0^2 \sin^2 \phi_1 \\ &= \Delta^2 k_0 \phi_1 - \sin^2 \phi_1 - \cos^2 2\phi; \end{aligned}$$

whence

$$\cos^2 2\phi + 2k_0 \sin^2 \phi_1 \cos 2\phi = \Delta^2 k_0 \phi_1 - \sin^2 \phi_1;$$

and from this,

$$\begin{aligned} \cos 2\phi &= -k_0 \sin^2 \phi_1 \pm \sqrt{\Delta^2 k_0 \phi_1 - \sin^2 \phi_1 + k_0^2 \sin^4 \phi_1} \\ (19) \quad &= \cos \phi_1 \Delta k_0 \phi_1 - k_0 \sin^2 \phi_1; \end{aligned}$$

whence, also,

$$\begin{aligned} 1 - \cos^2 2\phi &= 1 - \cos^2 \phi_1 \Delta^2 \phi_1 + 2k \sin^2 \phi_1 \cos \phi_1 \Delta k_0 \phi_1 - k_0^2 \sin^4 \phi_1 \\ &= \sin^2 \phi_1 (1 + k_0^2 \cos^2 \phi_1 + 2k_0 \cos \phi_1 \Delta k_0 \phi_1 - k_0^2 \sin^2 \phi_1) \end{aligned}$$

and

$$(20) \quad \sin 2\phi = \sin \phi_1 (\Delta k_0 \phi_1 + k_0 \cos \phi_1).$$

Differentiating equation (19), we get

$$2 \sin 2\phi \frac{d\phi}{d\phi_1} = \sin \phi_1 \frac{(k_0 \cos \phi_1 + \Delta k_0 \phi_1)^2}{\Delta k_0 \phi_1}.$$

Dividing this by equation (20), we have

$$\frac{2d\phi}{d\phi_1} = \frac{k_0 \cos \phi_1 + \Delta k_0 \phi_1}{\Delta k_0 \phi_1}.$$

But from (19), and eq. (6), Chap. IV,

$$\begin{aligned} k^2 \sin^2 \phi &= \frac{k^2 (1 - \cos 2\phi)}{2} \\ &= \frac{2k_0}{(1 + k_0)^2} \{1 + k_0 \sin^2 \phi_1 - \cos \phi_1 \Delta k_0 \phi_1\}; \end{aligned}$$

whence

$$\Delta k \phi = \frac{\Delta k_0 \phi_1 + k_0 \cos \phi_1}{1 + k_0},$$

and

$$2\Delta k \phi \cdot \frac{d\phi}{d\phi_1} = \frac{(k_0 \cos \phi_1 + \Delta k_0 \phi_1)^2}{(1 + k_0) \Delta k_0 \phi_1},$$

and

$$d\phi \Delta k \phi = \frac{d\phi_1}{\Delta k_0 \phi_1} \cdot \frac{(k_0 \cos \phi_1 + \Delta k_0 \phi_1)^2}{2(1 + k_0)}.$$

This gives immediately, by integration,

$$\begin{aligned} Ek\phi &= \frac{1}{2(1 + k_0)} \int \frac{d\phi_1}{\Delta k_0 \phi_1} \{k_0 \cos \phi_1 + \Delta k_0 \phi_1\}^2 \\ &= \frac{1}{2(1 + k_0)} \int \frac{d\phi_1}{\Delta k_0 \phi_1} \{2\Delta^2 k_0 \phi_1 + 2k_0 \cos \phi_1 \Delta k_0 \phi_1 - k_1'^2\} \\ (21) \quad &= \frac{Ek_0 \phi_1}{1 + k_0} + \frac{k_0 \sin \phi_1}{1 + k_0} - \frac{1}{2}(1 - k_0) Fk_0 \phi_1. \end{aligned}$$

Thus the value of $Ek\phi$ is made to depend upon $Ek_0 \phi_1$ (containing a smaller modulus and a larger amplitude), and upon the integral of the first class, $Fk_0 \phi_1$; k_0 , ϕ_1 , etc., being determined by the formulæ (6) to (12) of Chap. IV.

By successive applications of equation (21), $Ek\phi$ may be made to depend ultimately upon $Ek_{0n} \phi_n$, where k_{0n} approximates to zero and $Ek_{0n} \phi_n$ to ϕ_n .

Or, by reversing, it may be made to depend upon $Ek_n \phi_{0n}$, where k_n approximates to unity and $Ek_n \phi_{0n}$ to $-\cos \phi_{0n}$.

To facilitate this, assume

$$Gk\phi = Ek\phi - Fk\phi.$$

Subtracting from equation (21) the equation

$$Fk\phi = \frac{1+k_0}{2} Fk_0\phi_1 \text{ (see eq. (13), Chap. IV),}$$

we have

$$Gk\phi = \frac{1}{1+k_0} (Gk_0\phi_1 + k_0 \sin \phi_1 - k_0 Fk_0\phi_1).$$

Repeated applications of this give

$$Gk_0\phi_1 = \frac{1}{1+k_{00}} (Gk_{00}\phi_2 + k_{00} \sin \phi_2 - k_{00} Fk_{00}\phi_2),$$

$$\cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot$$

$$Gk_{0(n-1)}\phi_{n-1} = \frac{1}{1+k_{0n}} (Gk_{0n}\phi_n + k_{0n} \sin \phi_n - k_{0n} Fk_{0n}\phi_n).$$

Whence

$$(22) \quad Gk\phi = \sum_n \left\{ \frac{k_{0n}(\sin \phi_n - Fk_{0n}\phi_n)}{[\frac{1}{II}]_n (1+k_{0n})} \right\} + \frac{Gk_{0n}\phi_n}{[\frac{1}{II}]_n (1+k_{0n})}.$$

But since (compare eq. (13), Chap. IV)

$$Fk\phi = \frac{Fk_{0n}\phi_n [\frac{1}{II}]_n (1+k_{0n})}{2^n},$$

or

$$(23) \quad \frac{Fk_{0n}\phi_n}{[\frac{1}{II}]_n (1+k_{0n})} = \frac{2^n Fk\phi}{[\frac{1}{II}]_n (1+k_{0n})^2};$$

and since, also, (compare eq. (6), Chap. IV,)

$$\frac{k_{0(n-1)}^2}{k_{0n}} = \frac{2^2}{(1 + k_{0n})^2},$$

we have

$$\begin{aligned} (24) \quad \frac{2^n k_{0n}}{[\text{II}]_n (1 + k_{0n})^2} &= \frac{k_{0n}}{2^n} \left[\frac{\text{I}}{\text{II}} \right]_n \frac{k_{0(n-1)}^2}{k_{0n}} \\ &= \frac{k_{0n}}{2^n} \left[\frac{\text{I}}{\text{II}} \right]_n \frac{k_{0(n-1)}}{k_{0n}} \left[\frac{\text{I}}{\text{II}} \right]_n k_{0(n-1)} \\ &= \frac{k_{0n}}{2^n} \cdot \frac{k}{k_0} \cdot \frac{k_0}{k_{00}} \cdot \dots \cdot \frac{k_{0(n-1)}}{k_{0n}} \cdot k \left[\frac{\text{I}}{\text{II}} \right]_n k_{0(n-1)} \\ &= \frac{k^2}{2^n} \left[\frac{\text{I}}{\text{II}} \right]_n k_{0(n-1)}. \end{aligned}$$

Substituting these values in equation (22), and neglecting the term containing $Gk_{0n}\phi_n$ since, carried to its limiting value,

$$\begin{aligned} Gk_{0n}\phi_n &= Ek_{0n}\phi_n - Fk_{0n}\phi_n \\ &= \phi_n - \phi_n = 0, \end{aligned} \quad (n = \text{limiting value,})$$

we have

$$\begin{aligned} (25) \quad Gk\phi &= \sum_n \left\{ \frac{k \sqrt{k_{0n}} \sin \phi_n [\text{II}]_n \sqrt{k_{0(n-1)}} - k^2 [\text{I}]_n k_{0(n-1)}}{2^n} \right\} \\ &= k \left[\frac{\sqrt{k_0}}{2} \sin \phi_1 + \frac{\sqrt{k_0 k_{00}}}{2^2} \sin \phi_2 + \frac{\sqrt{k_0 k_{00} k_{003}}}{2^3} \sin \phi_3 + \dots \right] \\ &\quad - \frac{k^2}{2} \left[1 + \frac{k_0}{2} + \frac{k_0 k_{00}}{2^2} + \frac{k_0 k_{00} k_{003}}{2^3} + \dots \right]; \end{aligned}$$

whence

$$\begin{aligned} (26) \quad Ek\phi &= Fk\phi \left[1 - \frac{k^2}{2} \left(1 + \frac{k_0}{2} + \frac{k_0 k_{00}}{2^2} + \dots \right) \right] \\ &+ k \left[\frac{\sqrt{k_0}}{2} \sin \phi_1 + \frac{\sqrt{k_0 k_{00}}}{2^2} \sin \phi_2 + \frac{\sqrt{k_0 k_{00} k_{003}}}{2^3} \sin \phi_3 + \dots \right]. \end{aligned}$$

From eq. (3), Chap. V, we see that when $\phi = \frac{\pi}{2}$,

$$\phi_n = 2^{n-1}\pi.$$

Substituting these values in equation (26), we have for a complete Elliptic Integral of the second class,

$$(27) \quad E\left(k, \frac{\pi}{2}\right) = F\left(k, \frac{\pi}{2}\right) \left[1 - \frac{k^2}{2} \left(1 + \frac{k_0}{2} + \frac{k_0 k_{00}}{2^2} + \frac{k_0 k_{00} k_{03}}{2^3} + \dots \right) \right].$$

In a similar manner we could have found the formula for $E(k, \phi)$ in terms of an increasing modulus, viz.,

$$(28) \quad E(k, \phi) = F(k, \phi) \left[1 + k \left(1 + \frac{2}{k_1} + \frac{2^2}{k_1 k_2} + \frac{2^3}{k_1 k_2 k_3} + \dots \right. \right. \\ \left. \left. + \frac{2^{n-2}}{k_1 k_2 \dots k_{n-2}} - \frac{2^{n-1}}{k_1 k_2 \dots k_{n-1}} \right) \right] \\ - k \left[\sin \phi + \frac{2}{\sqrt{k}} \sin \phi_1 + \frac{2^2}{\sqrt{k k_1}} \sin \phi_2 + \dots \right. \\ \left. + \frac{2^{n-1}}{\sqrt{k k_1 \dots k_{n-2}}} \sin \phi_{n-1} - \frac{2^n}{\sqrt{k k_1 \dots k_{n-1}}} \sin \phi_n \right].$$

CHAPTER XI.

ELLIPTIC INTEGRALS OF THE THIRD ORDER.

THE Elliptic Integral of the third order is

$$(1) \quad \Pi(n, k, \phi) = \int_0^\phi \frac{d\phi}{(1 + n \sin^2 \phi) \Delta \phi}.$$

Put

$$(2) \quad \Pi(\phi) + \Pi(\psi) = S;$$

whence we have immediately

$$(3) \quad dS = \frac{d\phi}{(1 + n \sin^2 \phi) \Delta \phi} + \frac{d\psi}{(1 + n \sin^2 \psi) \Delta \psi}.$$

But, eq. (2), Chap. II,

$$(4) \quad \frac{d\phi}{\Delta \phi} + \frac{d\psi}{\Delta \psi} = 0;$$

whence

$$(5) \quad \begin{aligned} dS &= \left(\frac{1}{1 + n \sin^2 \phi} - \frac{1}{1 + n \sin^2 \psi} \right) \frac{d\phi}{\Delta \phi} \\ &= \frac{n(\sin^2 \psi - \sin^2 \phi)}{(1 + n \sin^2 \phi)(1 + n \sin^2 \psi)} \cdot \frac{d\phi}{\Delta \phi}. \end{aligned}$$

From equation (8), Chap. X, we get by differentiation, since σ (or μ) is constant,

$$\Delta \phi \cdot d\phi + \Delta \psi \cdot d\psi = k^2 \sin \sigma d(\sin \phi \sin \psi),$$

or, from equation (3),

$$(\sin^2 \psi - \sin^2 \phi) \frac{d\phi}{\Delta \phi} = \sin \sigma d(\sin \phi \sin \psi).$$

This, introduced into equation (5), gives

$$dS = \frac{n \sin \sigma d(\sin \phi \sin \psi)}{1 + n (\sin^2 \phi + \sin^2 \psi) + n^2 \sin^2 \phi \sin^2 \psi}.$$

Put

$$\sin \phi \sin \psi = q, \quad \sin^2 \phi + \sin^2 \psi = p;$$

whence

$$(6) \quad dS = \frac{n \sin \sigma dq}{1 + np + n^2 p^2}.$$

From equation (5), Chap. II, we have

$$\cos \sigma = \cos \phi \cos \psi - \sin \phi \sin \psi \Delta \sigma,$$

from which we easily get

$$\begin{aligned} (\cos \sigma + q \Delta \sigma)^2 &= \cos^2 \phi \cos^2 \psi \\ &= (1 - \sin^2 \phi)(1 - \sin^2 \psi) \\ &= 1 - p + q^2, \end{aligned}$$

and thence

$$\begin{aligned} p &= 1 + q^2 - (\cos \sigma + q \Delta \sigma)^2 \\ &= \sin^2 \sigma - 2 \cos \sigma \Delta \sigma q + k^2 \sin^2 \sigma \cdot q^2. \end{aligned}$$

This, substituted in eq. (6), gives

$$\begin{aligned} dS &= \frac{n \sin \sigma dq}{1 + n \sin^2 \sigma - 2n \cos \sigma \Delta \sigma q + n(n + k^2 \sin^2 \sigma)q^2} \\ &= \frac{n \sin \sigma dq}{A - 2Bq + Cq^2}, \end{aligned}$$

where

$$\begin{aligned} A &= 1 + n \sin^2 \sigma, \\ B &= n \cos \sigma \Delta \sigma, \\ C &= nk^2 \sin^2 \sigma + n^2. \end{aligned}$$

From this we get

$$S = n \sin \sigma \int \frac{dq}{A - 2Bq + Cq^2} + \text{Const.}$$

In order to determine the constant of integration we must observe that for $\phi = 0$, $\psi = \sigma$ and $q = 0$; whence

$$\Pi\sigma = n \sin \sigma \int_{q=0} \frac{dq}{A - 2Bq + Cq^2} + \text{Const.};$$

whence

$$S = \Pi\sigma + n \sin \sigma \int_0^q \frac{dq}{A - 2Bq + Cq^2},$$

or

$$(7) \quad \Pi\phi + \Pi\psi = \Pi\sigma + n \sin \sigma \int_0^q \frac{dq}{A - 2Bq + Cq^2}.$$

But we have

$$\begin{aligned} dS &= \frac{CMdq}{AC - B^2 + (Cq - B)^2} \\ &= \frac{CM}{AC - B^2} \cdot \frac{dq}{1 + \left(\frac{Cq - B}{\sqrt{AC - B^2}}\right)^2} \\ &= \frac{M}{\sqrt{AC - B^2}} \cdot \frac{\frac{Cdq}{\sqrt{AC - B^2}}}{1 + \left(\frac{Cq - B}{\sqrt{AC - B^2}}\right)^2} \end{aligned}$$

where $M = n \sin \sigma$.

The integral of the second member is

$$\frac{M}{\sqrt{AC - B^2}} \tan^{-1} \frac{Cq - B}{\sqrt{AC - B^2}};$$

whence

$$\int_0^q dS = S_1 = \frac{M}{\sqrt{AC-B^2}} \left[\tan^{-1} \frac{Cq-B}{\sqrt{AC-B^2}} + \tan^{-1} \frac{B}{\sqrt{AC-B^2}} \right];$$

or, since

$$\tan^{-1} x + \tan^{-1} y = \tan^{-1} \frac{x+y}{1-xy},$$

$$S_1 = \frac{M}{\sqrt{AC-B^2}} \tan^{-1} q \frac{\sqrt{AC-B^2}}{A-Bq}.$$

Substituting the values of A , B , C and M , we have

$$\begin{aligned} AC-B^2 &= n(1+n-\Delta^2\sigma)(1+n\sin^2\sigma) - n^2\cos^2\sigma\Delta^2\sigma \\ &= n(1+n-\Delta^2\sigma + n(1+n)\sin^2\sigma - n\Delta^2\sigma) \\ &= n(1+n)(1-\Delta^2\sigma + n\sin^2\sigma) \\ &= n(1+n)(k^2+n)\sin^2\sigma; \end{aligned}$$

and putting

$$\frac{(1+n)(k^2+n)}{n} = \Omega,$$

we have

$$\sqrt{AC-B^2} = n\sqrt{\Omega}\sin\sigma.$$

Substituting these values in eq. (7), we have

$$\begin{aligned} \Pi(n, k, \phi) + \Pi(n, k, \psi) - \Pi(n, k, \sigma) &= S_1 \\ &= \frac{1}{\sqrt{\Omega}} \tan^{-1} \frac{n\sqrt{\Omega}\sin\phi\sin\psi\sin\sigma}{1+n\sin^2\sigma - n\sin\phi\sin\psi\cos\sigma\Delta\sigma}. \end{aligned}$$

CHAPTER XII.

NUMERICAL CALCULATIONS. q .

CALCULATION OF THE VALUE OF q .

FROM eq. (7), Chap. IX, we have

$$\operatorname{dn} u = \sqrt{k'} \frac{\Theta(u + K)}{\Theta(u)};$$

whence, eq. (9), Chap. IV, eqs. (27) and (39), Chap. IX,

$$\begin{aligned} (1) \quad \sqrt{\cos \theta} &= \frac{1 - 2q + 2q^4 - 2q^9 + 2q^{16} - \dots}{1 + 2q + 2q^4 + 2q^9 + 2q^{16} + \dots} \\ &= 1 - 4q + 8q^2 - 16q^3 + 32q^4 - 56q^5 + \dots \end{aligned}$$

The first five terms of this series can be represented by

$$\sqrt{\cos \theta} = \frac{1 - 2q}{1 + 2q}.$$

From this we get

$$(2) \quad q = \frac{1}{2} \cdot \frac{1 - \sqrt{\cos \theta}}{1 + \sqrt{\cos \theta}},$$

which is exact up to the term containing q^5 .

Or we can deduce a more exact formula as follows: From eq. (1),

$$\begin{aligned} \frac{1 + \sqrt{\cos \theta}}{1 - \sqrt{\cos \theta}} &= \frac{\sqrt{1 + \tan^2 \frac{1}{2}\theta} + \sqrt{1 - \tan^2 \frac{1}{2}\theta}}{\sqrt{1 + \tan^2 \frac{1}{2}\theta} - \sqrt{1 - \tan^2 \frac{1}{2}\theta}} \\ &= \frac{1 + 2q^4 + 2q^{16} + \dots}{2q + 2q^9 + 2q^{25} + \dots}; \end{aligned}$$

whence, by the method of indeterminate coefficients,

$$(3) \quad q = \frac{1}{4} \tan^2 \frac{\theta}{2} + \frac{1}{16} \tan^4 \frac{\theta}{2} + \frac{5}{512} \tan^6 \frac{\theta}{2} + \frac{45}{2648} \tan^8 \frac{\theta}{2} + \dots,$$

or

$$\log q = 2 \log \tan \frac{\theta}{2} - \log 4 +$$

$$\log \left(1 + \frac{1}{4} \tan^2 \frac{\theta}{2} + \frac{1}{128} \tan^4 \frac{\theta}{2} + \frac{45}{512} \tan^6 \frac{\theta}{2} + \dots \right)$$

$$(4) \quad = 2 \log \tan \frac{\theta}{2} - \log 4 +$$

$$M \left(\frac{1}{4} \tan^2 \frac{\theta}{2} + \frac{1}{128} \tan^4 \frac{\theta}{2} + \frac{45}{512} \tan^6 \frac{\theta}{2} + \dots \right),$$

M being the modulus of the common system of logarithms.

Put

$$(5) \quad \log q = 2 \log \tan \frac{\theta}{2} + 9.397940 + a \tan^2 \frac{\theta}{2} +$$

$$b \tan^4 \frac{\theta}{2} + c \tan^6 \frac{\theta}{2} + \dots,$$

in which

$$\log a = 9.0357243;$$

$$\log b = 8.64452;$$

$$\log c = 8.41518;$$

$$\log d = 8.25283.$$

EXAMPLE. Let $k' = \cos 10^\circ 23' 46''$. To find *q*.

$$4 \log \tan \frac{\theta}{2} = 5.835 \qquad 2 \log \tan \frac{\theta}{2} = 7.9176842$$

$$\log a = 9.036 \qquad 9.3979400$$

$$\hline 4.871 \qquad 74$$

$$\log q = 7.3156316$$

$$a \tan^2 \frac{\theta}{2} = 0.0000074$$

When θ approaches 90° , $\tan \frac{\theta}{2}$ differs little from unity, and the series in eq. (5) is not very converging, but q can be calculated by means of eq. (6), Chap. VII, viz.,

$$q = e^{-\frac{\pi K'}{K}}, \quad q' = e^{-\frac{\pi K}{K'}}.$$

By comparing these equations with eqs. (6) and (9), Chap. IV, we see that if

$$q = f(k) = f(\theta),$$

then

$$q' = f(k') = f(90^\circ - \theta).$$

Therefore, having θ , we can from its complement, $90^\circ - \theta$, find q' by eq. (5), and thence q by the following process. We have

$$\frac{1}{q} = e^{\frac{\pi K'}{K}}, \quad \frac{1}{q'} = e^{\frac{\pi K}{K'}};$$

whence

$$\log \frac{1}{q} \log \frac{1}{q'} = M^2 \pi^2 = 1.8615228,$$

$$(6) \quad \log \log \frac{1}{q} + \log \log \frac{1}{q'} = 0.2698684,$$

by which we can deduce q from q' .

EXAMPLE. Let $\theta = 79^\circ 36' 14''$. To find q .

$$90^\circ - \theta = 10^\circ 23' 46''.$$

By eq. (5) we get

$$\log q' = 7.3156316, \quad \log \frac{1}{q'} = 2.6843684,$$

$$\text{and } \log \log \frac{1}{q'} = .4288421;$$

and by eq. (6),

$$\log \log \frac{1}{q} = 9.8410263;$$

whence

$$\log q = \bar{1}.3065321.$$

When $k' = k = \cos 45^\circ = \frac{1}{2}\sqrt{2}$, eq. (6) becomes

$$(7) \quad \log \frac{1}{q} = M\pi = 1.3643763; \quad (k = k';)$$

whence

$$\begin{aligned} \log q &= 2.6356237, \\ q &= 0.0432138. \end{aligned} \quad (k = k'.)$$

EXAMPLE. Given $\theta = 10^\circ 23' 46''$. Find q .

$$\text{Ans. } \log q = 7.3156316.$$

EXAMPLE. Given $\theta = 82^\circ 45'$. Find q .

$$\text{Ans. } \log q = 9.37919.$$

CHAPTER XIII.

NUMERICAL CALCULATIONS. K .

CALCULATION OF THE VALUE OF K .

We have already found from eq. (37), Chap. IX,

$$(1) \quad \Theta(0) = \sqrt{\frac{2k'K}{\pi}},$$

and from eq. (40), same chapter,

$$(2) \quad \Theta(K) = \frac{\Theta(0)}{\sqrt{k'}} = \sqrt{\frac{2K}{\pi}}.$$

But, eqs. (38) and (27), Chap. IX,

$$\Theta(K) = 1 + 2q + 2q^4 + 2q^9 + 2q^{16} + \dots,$$

$$\Theta(0) = 1 - 2q + 2q^4 - 2q^9 + 2q^{16} - \dots;$$

whence, eq. (2),

$$(3) \quad K = \frac{\pi}{2} (1 + 2q + 2q^4 + 2q^9 + \dots)^2.$$

By adding eqs. (1) and (2) we get

$$\Theta(0) + \Theta(K) = \sqrt{\frac{2K}{\pi}} (1 + \sqrt{k'});$$

whence

$$\begin{aligned} K &= \frac{\pi}{2} \left(\frac{\Theta(0) + \Theta(K)}{1 + \sqrt{k'}} \right)^2 \\ &= \frac{\pi}{2} \left[\frac{2(1 + 2q + 2q^4 + 2q^9 + \dots)}{1 + \sqrt{k'}} \right]^2 \end{aligned}$$

$$(4) \quad = \frac{\pi}{2} \left(\frac{2}{1 + \sqrt{k'}} \right)^2 (1 + 2q^4 + 2q^{16} + \dots)^2.$$

EXAMPLE. Let $k = \sin \theta = \sin 19^\circ 30'$. Required K .

First Method. By eq. (3).

By eq. (5), Chap. XII, we find $\log q = 8.6356236$. Applying eq. (3), using only two terms of the series, we have

$$\begin{aligned} 1 + 2q &= 1.0147662 \\ \log(1 + 2q) &= 0.0063660 \\ 2 \log(1 + 2q) &= 0.0127320 \\ \log \frac{\pi}{2} &= 0.1961199 \\ \hline \log K &= 0.2088519 \\ K &= 1.615101 \end{aligned}$$

Second Method. By eq. (4).

Equation (4) may be written, neglecting q^4 ,

$$K = \frac{\pi}{2} \left(\frac{1 + \sqrt{\cos \theta}}{2} \right)^{-2};$$

whence

$$\begin{aligned} \log \cos \theta &= 9.9743466, \\ \log \sqrt{\cos \theta} &= 9.9871733, \\ 1 + \sqrt{\cos \theta} &= 1.9708973, \\ \frac{1 + \sqrt{\cos \theta}}{2} &= 0.98544865; \end{aligned}$$

and

$$\begin{aligned} \log K &= 0.2088519, \\ K &= 1.615101, \end{aligned}$$

the same result as above.

Third Method. By eq. (7), Chap. V.

$\theta = 19^\circ 30'$	$\theta_0 = 1^\circ 41' 31''.1$
$\frac{1}{2}\theta = 9^\circ 45'$	$\frac{1}{2}\theta_0 = 0^\circ 50' 45''.5$
$\log \tan \frac{1}{2}\theta = 9.235103$	
$\log \cos \frac{1}{2}\theta = 9.993681$	$\log \cos \frac{1}{2}\theta_0 = 9.999953$
$\left. \begin{array}{l} \log \tan^2 \frac{1}{2}\theta \\ \log \sin \theta_0 \end{array} \right\} = 8.470206$	
$\theta_0 = 1^\circ 41' 31''.1$	

$$\begin{aligned} \log \cos^2 \frac{1}{2}\theta &= 9.987362 \\ \log \cos^2 \frac{1}{2}\theta_0 &= 9.999906 \end{aligned}$$

$$9.987268$$

$$\log \frac{\pi}{2} = 0.196120$$

$$\log K = 0.208852$$

θ_{00} is not calculated, as it is evident that its cosine will be 1.

EXAMPLE. Given $k = \sin 75^\circ$. Find K .

By eq. (7), Chap. V.

From eqs. (14), Chap. IV, we find

$$k = \sin \theta = \sin 75^\circ \qquad \log = 9.9849438$$

$$k_0 = \left\{ \begin{array}{l} \tan^2 \frac{1}{2}\theta = \tan^2 37^\circ 30' \\ \sin \theta_0 = \sin 36^\circ 4' 16''.47 \end{array} \right\} \qquad 9.7699610$$

$$k_{00} = \left\{ \begin{array}{l} \tan^2 \frac{1}{2}\theta_0 = \tan^2 18^\circ 2' 8''.235 \\ \sin \theta_{00} = \sin 6^\circ 5' 9''.38 \end{array} \right\} \qquad 9.0253880$$

$$k_{03} = \left\{ \begin{array}{l} \tan^2 \frac{1}{2}\theta_{00} = \tan^2 3^\circ 2' 34''.69 \\ \sin \theta_{03} = \sin 9' 42''.90 \end{array} \right\} \qquad 7.4511672$$

	log	2 log	a. c. 2 log
$\cos \frac{1}{2}\theta = \cos 37^\circ 30'$	9.8994667	9.7989334	0.2010666
$\cos \frac{1}{2}\theta_0 = \cos 18^\circ 2'.13725$	9.9781184	9.9562368	0.0437632
$\cos \frac{1}{2}\theta_{02} = \cos 3^\circ 2'.57817$	9.9993873	9.9987746	0.0012254
$\cos \frac{1}{2}\theta_{03} = \cos 4'.8575$	9.9999995	9.9999990	0.0000010

0.2460562

$\frac{\pi}{2}$

$\frac{\pi}{2}$

.1961199

log $K =$ 0.4421761

$K = 2.768064$ *Ans.*

EXAMPLE. Given $k = \sin 45^\circ$. Find K .

Method of eq. (7), Chap. V.

From eqs. (14), Chap. IV, we have

	log
$k_0 = \left\{ \begin{array}{l} \tan^2 \frac{1}{2}\theta = \tan^2 22^\circ 30' \\ \sin \theta_0 = \sin 9^\circ 52'.75683 \end{array} \right\}$	9.2344486
$k_{00} = \left\{ \begin{array}{l} \tan^2 \frac{1}{2}\theta_0 = \tan^2 4^\circ 56'.37841 \\ \sin \theta_{00} = \sin 25'.679 \end{array} \right\}$	7.8733009
$k_{03} = \left\{ \begin{array}{l} \tan^2 \frac{1}{2}\theta_{00} = \tan^2 12'.3395 \\ \sin \theta_{03} = \sin 0'.05 \end{array} \right\}$	5.1445523

a. c. log $\cos^2 \frac{1}{2}\theta$ 0.0687694

a. c. log $\cos^2 \frac{1}{2}\theta_0$ 0.0032320

a. c. log $\cos^2 \frac{1}{2}\theta_{00}$ 0.0000060

log $\frac{\pi}{2}$ 0.1961199

log $K =$ 0.2681273

$K = 1.8540747$ *Ans.*

EXAMPLE. Given $\theta = 63^\circ 30'$. Find K .

Ans. log $K = 0.3539686$.

EXAMPLE. Given $\theta = 34^\circ 30'$. Find K .

Ans. $K = 1.72627$.

CHAPTER XIV.

NUMERICAL CALCULATIONS. *u.*

CALCULATION OF THE VALUE OF *u.*

WHEN $\theta^\circ = \sin^{-1}k < 45^\circ$.

EXAMPLE. Let $\phi = 30^\circ$, $k = \sin 45^\circ$. Find *u*.

First Method. Eq. (23), Chap. IV, and eqs. (14₁), (14₂), (14₃), Chap. IV.

By equations (14₁),

$$\frac{\theta}{2} = 22^\circ 30' ;$$

$$\log \tan \frac{\theta}{2} = 9.6172243 ;$$

$$\log \tan^2 \frac{\theta}{2} = 9.2344486 = \log k_0 = \log \sin \theta_0 ;$$

$$\theta_0 = 9^\circ 52' 45'' .41 ;$$

$$\log \tan \frac{\theta_0}{2} = 8.9366506 ;$$

$$\log \tan^2 \frac{\theta_0}{2} = 7.8733012 = \log k_{00} = \log \sin \theta_{00} ;$$

$$\theta_{00} = 0^\circ 25' 40'' .7 ;$$

$$\log \tan^2 \frac{\theta_{00}}{2} = 5.144552 = \log k_{00} .$$

By equations (14₂),

$$\begin{aligned}\phi &= 30^\circ \\ \log \tan \phi &= 9.761439 \\ \log \cos \theta &= 9.849485 \\ \hline \log \tan (\phi_1 - \phi) &= 9.610924 \\ \phi_1 - \phi &= 22^\circ 12' 27''.56 \\ \phi_1 &= 52^\circ 12' 27''.56\end{aligned}$$

$$\begin{aligned}\log \tan \phi_1 &= 0.110438 \\ \log \cos \theta_0 &= 9.993512 \\ \hline \log \tan (\phi_2 - \phi_1) &= 0.103949 \\ \phi_2 - \phi_1 &= 51^\circ 47' 32''.59\end{aligned}$$

$$\begin{aligned}\phi_2 &= 104^\circ 0' 0''.15 \\ \log \tan \phi_2 &= 0.603228 \\ \log \cos \theta_{00} &= 9.999988 \\ \hline \log \tan (\phi_3 - \phi_2) &= 0.603216 \\ \phi_3 - \phi_2 &= 104^\circ 0' 1''.5\end{aligned}$$

$$\phi_3 = 208^\circ 0' 1''.65$$

Since $\frac{\phi_2}{4} = 26^\circ 0' 0''.04$ and $\frac{\phi_3}{8} = 26^\circ 0' 0''.21$,

we need not calculate ϕ_4 .

$$\frac{\phi_3}{8} = 93600''.21.$$

Reducing this to radians, we have

$$\log \frac{\phi_3}{8} = 9.656852.$$

Substituting in eq. (23), Chap. IV, we have, since $\cos \theta_{03} = 1$,

$$\text{a. c. } \log \cos \theta = 0.150515$$

$$\log \cos \theta_0 = 9.993512$$

$$\log \cos \theta_{00} = 9.999988$$

$$\hline 0.144014$$

$$0.072007 = \log \sqrt{\frac{\cos \theta_0 \cos \theta_{00}}{\cos \theta}}$$

$$\log \frac{\phi_3}{8} = 9.656852$$

$$\log u = 9.728859$$

$$u = 0.535623, \quad \text{Ans.}$$

When $\theta = \sin^{-1} k > 45^\circ$.

EXAMPLE. Given $k = \sin 75^\circ$, $\tan \phi = \sqrt{\frac{2}{\sqrt{3}}}$. To find

$F(k, \phi)$.

First Method. Bisected Amplitudes.

By equations (24) and (25), Chap. IV, we get

$$\phi = 47^\circ 3' 30''.91,$$

$$\phi_{\frac{1}{2}} = 25^\circ 36' 5''.64,$$

$$\phi_{\frac{1}{4}} = 13^\circ 6' 30''.98,$$

$$\phi_{\frac{1}{8}} = 6^\circ 35' 40''.74,$$

$$\phi_{\frac{1}{16}} = 3^\circ 18' 8''.75,$$

$$\phi_{\frac{1}{32}} = 1^\circ 39' 7''.43,$$

$$\beta = 45^\circ;$$

$$\beta_0 = 24^\circ 40' 10''.94;$$

$$\beta_{00} = 12^\circ 39' 15''.83;$$

$$\beta_{03} = 6^\circ 22' 8''.40;$$

$$\beta_{04} =$$

Substituting in equation (26), Chap. IV, we have

$$F(k, \phi) = 32 \times 1^\circ 39' 7''.43$$

$$= 52^\circ 51' 58''.03$$

$$= 0.9226878. \quad \text{Ans.}$$

Second Method. Equation (29), Chap. IV.

From equations (18_s), Chap. IV, we have

			log
$k = \cos \eta$	$= \cos 15^\circ 0' 0''.00$		9.9849438
$k' = \sin \eta$	$= \sin 15^\circ 0' 0''.00$		9.4129962
$k'_0 = \left\{ \begin{array}{l} \tan^2 \frac{1}{2} \eta = \tan^2 7^\circ 30' 0''.00 \\ \sin \eta_0 = \sin 0^\circ 59' 35''.25 \end{array} \right\}$			8.2388582
$k_1 = \cos \eta_0$	$= \cos 0^\circ 59' 35''.25$		9.9999348
$k'_{00} = \left\{ \begin{array}{l} \tan^2 \frac{1}{2} \eta_0 = \tan^2 0^\circ 29' 47''.62 \\ \sin \eta_{00} = \sin 0^\circ 0' 15''.49 \end{array} \right\}$			5.8757219
$k_2 = \cos \eta_{00}$	$= \cos 0^\circ 0' 15''.49$		0.0000000
$k'_{03} = (\frac{1}{2} k'_{00})^2$			1.1493838

From equations (18_s), Chap. IV, we get

$$\begin{aligned}
 \phi &= 47^\circ 3' 30''.95; \\
 2\phi_0 - \phi &= 45^\circ; \\
 \phi_0 &= 46^\circ 1' 45''.475; \\
 \phi_{02} &= 46^\circ 1' 29''.41; \\
 \phi_{03} &= 46^\circ 1' 29''.41; \\
 45^\circ + \frac{1}{2} \phi_3 &= 68^\circ 0' 44''.705.
 \end{aligned}$$

Substituting these values in eq. (29), Chap. IV, we get

$$\begin{aligned}
 F(k, \phi) &= \sqrt{\frac{k_1}{k}} \cdot \frac{1}{M} \cdot \log \tan 68^\circ 0' 44''.705 \\
 &= 0.9226877. \quad \text{Ans.}
 \end{aligned}$$

Third Method. Equation (23)*, Chap. IV.

From equations (14₁), Chap. IV, we have

$$\begin{aligned}
 k &= \sin \theta &= \sin 75^\circ & \text{ } o' \text{ } o'' & \log = 9.9849438 \\
 k' &= \cos \theta &= \cos 75^\circ & & 9.4129962 \\
 k_0 &= \left\{ \begin{array}{l} \tan^2 \frac{1}{2} \theta = \tan^2 37^\circ 30' \\ \sin \theta_0 = \sin 36^\circ 4' 16''.47 \end{array} \right\} & & & 9.7699610 \\
 k'_1 &= \cos \theta_0 & & & 9.9075648 \\
 k_{02} &= \left\{ \begin{array}{l} \tan^2 \frac{1}{2} \theta_0 = \tan^2 18^\circ 2' 8''.235 \\ \sin \theta_{00} = \sin 6^\circ 5' 9''.38 \end{array} \right\} & & & 9.0253880 \\
 k'_2 &= \cos \theta_{00} & & & 9.9975452 \\
 k_{03} &= \left\{ \begin{array}{l} \tan^2 \frac{1}{2} \theta_{00} = \tan^2 3^\circ 2' 34''.69 \\ \sin \theta_{03} = \sin 9' 42''.90 \end{array} \right\} & & & 7.4511672 \\
 k'_3 &= \cos \theta_{03} & & & 9.9999982 \\
 k_{04} &= \left(\frac{1}{2} k_{03} \right)^2 & & & 4.3002761 \\
 k'_4 &= & & & 0.0000000
 \end{aligned}$$

From equations (14₂), Chap. IV, we have

$$\begin{aligned}
 \phi &= 47^\circ 3' 30''.94; \\
 \phi_1 &= 62^\circ 36' 3''.10; \\
 \phi_2 &= 119^\circ 55' 47''.67; \\
 \phi_3 &= 240^\circ o' o''.19; \\
 \phi_4 &= 480^\circ o' o''.
 \end{aligned}$$

Therefore the limit of $\phi, \frac{\phi_1}{2}, \frac{\phi_2}{4},$ or $\frac{\phi_n}{2^n}$ is $30^\circ = \frac{\pi}{6}$.

Substituting these values in eq. (23)*, Chap. IV, we have

$$\begin{aligned}
 F(k, \phi) &= \sqrt{\frac{k'_1 k'_2 k'_3 k'_4}{k'}} \cdot \frac{\pi}{6} \\
 &= 0.9226874. \quad \text{Ans.}
 \end{aligned}$$

EXAMPLE. Given $\phi = 30^\circ, k = \sin 89^\circ$. Find u .

Method of eq. (28), Chap. IV.

From eqs. (18₁) we find

$$k_1 = \sin \theta_1 \quad \text{and} \quad \tan^2 \frac{1}{2} \theta_1 = k = \sin \theta,$$

from which we find that $k_1 = 1$ as far as seven decimal places.

From eqs. (18₂) we have

$$\begin{aligned} \sin \phi &= 9.6989700 \\ k &= 9.9999338 \\ \sin (2\phi_0 - \phi) &= 9.6989038 \\ 2\phi_0 - \phi &= 29^\circ 59'.69733 \\ 2\phi_0 &= 59^\circ 59'.69733 \\ 45^\circ + \frac{1}{2} \phi_0^* &= 59^\circ 59'.92433 \\ \log (45^\circ + \frac{1}{2} \phi_0) &= 0.2385385 \end{aligned}$$

From eqs. (18₃), Chap. IV, we have

$$k = \cos \eta = \cos 1^\circ, \quad \frac{1}{2} \eta = 30'.$$

Substituting in eq. (28), Chap. IV, we have

$$\begin{aligned} \text{a. c. } \log \cos \frac{1}{2} \eta & 0.0000330 \\ \log \log (45^\circ + \frac{1}{2} \phi_0) & 9.3775585 \\ \text{a. c. } \log M & 0.3622157 \\ \log F(k, \phi) &= 9.7398072 \\ F(k, \phi) &= 0.549297. \quad \text{Ans.} \end{aligned}$$

EXAMPLE. Given $\phi = 79^\circ$, $k = 0.25882$. Find u .

$$\text{Ans. } u = 0.39947.$$

EXAMPLE. Given $\phi = 37^\circ$, $k = 0.86603$. Find u .

$$\text{Ans. } u = 0.68141.$$

* Since $k_1 = 1$, $\phi_{00} = \phi_0$, and we need not carry the calculation further.

CHAPTER XV.

NUMERICAL CALCULATIONS. ϕ .

EXAMPLE. Given $u = 1.368407$, $\theta = 38^\circ$. Find ϕ .

First Method. From eqs. (46) and (41)*, Chap. IX, we have

$$u = x\Theta^2(K),$$

$$\Delta\phi = \sqrt{k'} \frac{\Theta_1(x)}{\Theta(x)}.$$

From equations (5), Chap. XII, and (38), Chap. IX, we have

$$\begin{aligned} \log q &= 8.4734187 \\ \log \Theta^2(K) &= 0.0501955 \\ \log u &= 0.1362153 \\ \hline \log x &= 0.0860198 \\ x &= 69^\circ 50' 46''.12 \end{aligned}$$

From equations (23) and (24), Chap. IX, we get

$$\begin{aligned} \log \Theta_1(x) &= 9.9798368 \\ \log \Theta(x) &= 0.0192687 \\ \hline &9.9605681 \\ \log \sqrt{k'} &= 9.9482661 \\ \hline \log \Delta\phi &= 9.9088342 = \log \sin \lambda \end{aligned}$$

But

$$\begin{aligned} k^3 \sin^2 \phi &= 1 - \Delta^2 \phi, \\ k \sin \phi &= \cos \lambda; \end{aligned}$$

whence

$$\log \cos \lambda = 9.7675483$$

$$\log k = 9.7893420$$

$$\log \sin \phi = 9.9782063$$

$$\phi = 72^\circ. \text{ Ans.}$$

Second Method. From eq. (1), Chap. VI.

From eqs. (14₁) Chap. IV, we find

$$\begin{aligned} k_0 &= \left\{ \begin{array}{l} \tan^2 \frac{1}{2} \theta = \tan^2 19^\circ \\ \sin \theta_0 = \sin 6^\circ 48' .54569 \end{array} \right\} \log = 9.0739438 \\ &\quad \cos \theta_0 \quad \quad \quad 9.9969260 \\ k_{00} &= \left\{ \begin{array}{l} \tan^2 \frac{1}{2} \theta_0 = \tan^2 3^\circ 24' .2784 \\ \sin \theta_{00} = 12' .16659 \end{array} \right\} \quad \quad 7.5488952 \\ &\quad \cos \theta_{00} \quad \quad \quad 9.9999974 \\ k_{03} &= \left\{ \begin{array}{l} \tan^2 \frac{1}{2} \theta_{00} = \tan^2 6' .08329 \\ \sin \theta_{03} \end{array} \right\} \quad \quad 4.4957316 \\ &\quad \cos \theta_{03} \quad \quad \quad 0.0000000 \end{aligned}$$

Substituting these values in eq. (1), Chap. VI, we have

$$\begin{aligned} &\log \cos \theta_0 \quad 9.9969260 \\ &\log \cos \theta_{00} \quad 9.9999974 \\ &\quad \quad \quad 9.9969234 \\ &\log \sqrt{\cos \theta_0 \cos \theta_{00}} \quad 9.9984617 \\ \text{a. c. log} \quad \quad \quad &\quad \quad 0.0015383 \\ &\log u \quad .1362153 \\ &\log \sqrt{\cos \theta} \quad 9.9482660 \\ &\log 2^3 \quad .9030900^* \\ \text{a. c. log} \sqrt{\cos \theta_0 \cos \theta_{00}} &\quad 0.0015383 \\ &\quad \quad \quad 0.9891096 \\ &\quad \quad \quad 2.2418773 \\ &\log \phi_3^* \quad 2.7472323 \\ &\phi_3 \quad 558^\circ 46' .140 \end{aligned}$$

* n is taken equal to 3, because $\cos_{03} = 1$.

Whence, by equations (1)* of Chap. VI, we get

$$\begin{array}{rcl}
 k_{00} \log & = & 4.4957316 \\
 \sin \phi_2 & & 9.5075232_n \\
 \hline
 \sin (2\phi_2 - \phi_1) & & 4.0032548_n \\
 2\phi_2 - \phi_1 & = & -0'.00346 \\
 \phi_2 & = & 279^\circ 23'.06827 \\
 \\
 k_{00} \log & = & 7.5488952 \\
 \sin \phi_2 & & 9.9941484_n \\
 \hline
 \sin (2\phi_1 - \phi_2) & & 7.5430436_n \\
 2\phi_1 - \phi_2 & = & -12'.0039 \\
 \phi_1 & = & 139^\circ 35'.5321 \\
 \\
 k_0 \log & = & 9.0739438 \\
 \sin \phi_1 & & 9.8117249 \\
 \hline
 \sin (2\phi - \phi_1) & & 8.8856687 \\
 2\phi - \phi_1 & = & 4^\circ 24'.467 \\
 \phi & = & 71^\circ 59'.9999 \\
 & = & 72^\circ. \text{ Ans.}
 \end{array}$$

EXAMPLE. Given $u = 2.41569$, $\theta = 80^\circ$. Find ϕ .

Ans. $\phi = 82^\circ$.

EXAMPLE. Given $u = 1.62530$, $k = \frac{1}{2}$. Find ϕ .

Ans. $\phi = 87^\circ$.

CHAPTER XVI.

NUMERICAL CALCULATIONS. $E(k, \phi)$.

First Method. By Chap. X, eqs. (15), (16), and (17).

EXAMPLE. Given $k = 0.9327$, $\phi = 80^\circ$. Find $E(k, \phi)$.

By eq. (15), Chap. X,

$$\begin{array}{ll}
 \phi = 80^\circ; & \gamma = 67^\circ 44'. ; \\
 \phi_{\frac{1}{2}} = 50^\circ 43'.6, & \gamma_{\frac{1}{2}} = 46^\circ 40'.4; \\
 \phi_{\frac{1}{4}} = 27^\circ 48'.5, & \gamma_{\frac{1}{4}} = 26^\circ 0'.1; \\
 \phi_{\frac{1}{8}} = 14^\circ 16'.7, & \gamma_{\frac{1}{8}} = 13^\circ 24'.0; \\
 \phi_{\frac{1}{16}} = 7^\circ 11'.3, & \gamma_{\frac{1}{16}} = 6^\circ 45'.2; \\
 \phi_{\frac{1}{32}} = 3^\circ 36'.0, & \log \sin \gamma_{\frac{1}{32}} = 8.77094; \\
 \phi_{\frac{1}{64}} = 0.062831. & \\
 \therefore \phi_{\frac{1}{64}}^8 < 0.0000001. &
 \end{array}$$

Whence, by eq. (17),

$$\begin{array}{rcl}
 E(k, \phi_{\frac{1}{32}}) & = & 0.062794 \\
 \sin \phi & \sin^2 \gamma_{\frac{1}{2}} & = 0.52116 \\
 2 \sin \phi_{\frac{1}{2}} & \sin^2 \gamma_{\frac{1}{4}} & = 0.29757 \\
 4 \sin \phi_{\frac{1}{4}} & \sin^2 \gamma_{\frac{1}{8}} & = 0.10023 \\
 8 \sin \phi_{\frac{1}{8}} & \sin^2 \gamma_{\frac{1}{16}} & = 0.02728 \\
 16 \sin \phi_{\frac{1}{16}} & \sin^2 \gamma_{\frac{1}{32}} & = 0.00697 \\
 & & \hline
 & & 0.95321
 \end{array}$$

Hence, by eq. (16),

$$\begin{aligned}
 E(k, \phi) &= 32E(k, \phi_{\frac{1}{32}}) - 0.95321 \\
 &= 2.0094 - 0.9532 = 1.0562.
 \end{aligned}$$

Second Method. By Chap. X, eq. (26).

EXAMPLE. Given $k = \sin 75^\circ$, $\tan \phi = \sqrt{\frac{2}{\sqrt{3}}}$. Find

$E(k, \phi)$.

From eqs. (14₁), Chap. IV, we have

$$k = \sin \theta = \sin 75^\circ 0' 0'' \quad \log = 9.9849438$$

$$k' = \cos \theta = \cos 75^\circ \quad 9.4129962$$

$$k_0 = \left\{ \begin{array}{l} \tan^2 \frac{1}{2}\theta = \tan^2 37^\circ 30' \\ \sin \theta_0 = \sin 36^\circ 4' 16''.47 \end{array} \right\} \quad 9.7699610$$

$$k_1' = \cos \theta_0 \quad 9.9075648$$

$$k_{02} = \left\{ \begin{array}{l} \tan^2 \frac{1}{2}\theta_0 = \tan^2 18^\circ 2' 8''.235 \\ \sin \theta_{00} = \sin 6^\circ 5' 9''.38 \end{array} \right\} \quad 9.0253880$$

$$k_2' = \cos \theta_{00} \quad 9.9975452$$

$$k_{03} = \left\{ \begin{array}{l} \tan^2 \frac{1}{2}\theta_{00} = \tan^2 3^\circ 2' 34''.69 \\ \sin \theta_{03} = \sin 9' 42''.90 \end{array} \right\} \quad 7.4511672$$

$$k_3' = \cos \theta_{03} \quad 9.9999982$$

$$k_{04} = (\frac{1}{2}k_{03})^2 \quad 4.3002761$$

$$k_4' = \quad 0.0000000$$

From eqs. (14₁), Chap. IV, we have

$$\phi = 47^\circ 3' 30''.94;$$

$$\phi_1 = 62^\circ 36' 3''.10;$$

$$\phi_2 = 119^\circ 55' 47''.67;$$

$$\phi_3 = 240^\circ 0' 0''.19.$$

Applying eq. (26), Chap. X, we have

k^2	$\log =$	9.9698876	
a. c. 2		9.6989700	
		9.6688576	.4665064
k_0		9.7699610	
a. c. 2		9.6989700	
		9.1377886	.1373373
k_{00}		9.0253880	
a. c. 2		9.6989700	
		7.8621466	.0072802
k_{03}		7.4511672	
a. c. 2		9.6989700	
		5.0132838	.0000103
			.6111342

$$1 - .6111342 = 0.3888658.$$

From eq. (23)*, Chap. IV, we find $F(k, \phi) = 0.9226874$.

Hence

$$F(k, \phi) \left[1 - \frac{k^2}{2} \left(1 + \frac{k_0}{2} + \dots \right) \right] = 0.3588016$$

$$\frac{k \sqrt{k_0}}{2} \sin \phi_1 = 0.3290186$$

$$\frac{k \sqrt{k_0 k_{00}}}{4} \sin \phi_2 = 0.0522872$$

$$\frac{k \sqrt{k_0 k_{02} k_{03}}}{8} \sin \phi_3 = -0.0013888$$

$$\frac{k \sqrt{k_0 \dots k_{04}}}{16} \sin \phi_4 = 0.0000010$$

$$0.3799180$$

Whence

$$E(k, \phi) = 0.3588016 + 0.3799180 = 0.7387196. \quad \text{Ans.}$$

EXAMPLE. Given $k = \sin 75^\circ$. Find $E\left(k, \frac{\pi}{2}\right)$.

From Example 2, Chap. XIII, we find

$$\log F\left(k, \frac{\pi}{2}\right) = 0.4421761$$

$$\log 0.3888658 = \underline{1.5897998}$$

$$\log E\left(k, \frac{\pi}{2}\right) = 0.0319759$$

$$E\left(k, \frac{\pi}{2}\right) = 1.076405. \quad \text{Ans.}$$

EXAMPLE. Given $k = \sin 30^\circ$, $\phi = 81^\circ$. Find $E(k, \phi)$.

$$\text{Ans. } E(k, \phi) = 1.33124.$$

EXAMPLE. Find $E(\sin 80^\circ, 55^\circ)$. Ans. 0.82417.

EXAMPLE. Find $E\left(\sin 27^\circ, \frac{\pi}{2}\right)$. Ans. 1.48642.

EXAMPLE. Find $E(\sin 19^\circ, 27^\circ)$. Ans. 0.46946.

CHAPTER XVII.

APPLICATIONS.

RECTIFICATION OF THE LEMNISCATE.

THE polar equation of the Lemniscate is $r = a \sqrt{\cos 2\theta}$, referred to the centre as the origin. From this we get

$$\frac{dr}{d\theta} = -\frac{a \sin 2\theta}{\sqrt{\cos 2\theta}};$$

whence the length of the arc measured from the vertex to any point whose co-ordinates are r and θ

$$\begin{aligned} s &= \int \left\{ \left(\frac{dr}{d\theta} \right)^2 + r^2 \right\}^{\frac{1}{2}} d\theta = a \int \left\{ \frac{\sin^2 2\theta}{\cos 2\theta} + \cos 2\theta \right\}^{\frac{1}{2}} d\theta \\ &= a \int \frac{d\theta}{\sqrt{\cos 2\theta}} = a \int \frac{d\theta}{\sqrt{1 - 2 \sin^2 \theta}}. \end{aligned}$$

Let $\cos 2\theta = \cos^2 \phi$, whence

$$\begin{aligned} s &= a \int \frac{\frac{d\theta}{d\phi} d\phi}{\cos \phi} = a \int \frac{\sin \phi d\phi}{\sqrt{1 - \cos^4 \phi}} \\ &= a \int_0^\phi \frac{d\phi}{\sqrt{1 + \cos^2 \phi}} = \frac{a}{\sqrt{2}} \int_0^\phi \frac{d\phi}{\sqrt{1 - \frac{1}{2} \sin^2 \phi}} \\ &= \frac{a}{\sqrt{2}} F\left(\frac{1}{\sqrt{2}}, \phi\right). \end{aligned}$$

Since $r = a \sqrt{\cos 2\theta} = a \cos \phi$, the angle ϕ can be easily constructed by describing upon the axis a of the Lemniscate a semicircle, and then revolving the radius vector until it cuts this semicircle. In the right-angled triangle of which this is one side, and the axis the hypotenuse, ϕ is evidently the angle between the axis and the revolved position of the radius vector.

RECTIFICATION OF THE ELLIPSE.

Since the equation of the ellipse is $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, we can assume $x = a \sin \phi$, $y = b \cos \phi$, so that ϕ is the complement of the *eccentric angle*. Hence

$$\begin{aligned} s &= \int \sqrt{dx^2 + dy^2} = a \int d\phi \sqrt{1 - e^2 \sin^2 \phi} \\ &= aE(e, \phi), \end{aligned}$$

in which e , the eccentricity of the ellipse, is the modulus of the Elliptic Integral.

The length of the Elliptic Quadrant is

$$s' = aE\left(e, \frac{\pi}{2}\right).$$

EXAMPLE. The equation of an ellipse is $\frac{x^2}{16.81} + \frac{y^2}{16} = 1$; required the length of an arc whose abscissas are 1.061162 and 4.100000 : of the quadrantal arc. *Ans.* 5.18912; 6.36189.

RECTIFICATION OF THE HYPERBOLA.

On the curve of the hyperbola, construct a straight line perpendicular to the axis x , and at a distance from the centre equal to the projection of b , the transverse axis, upon the asymptote, i.e. equal to $\frac{b^2}{\sqrt{a^2 + b^2}}$. Join the projection of the

given point of the hyperbola on this line with the centre. The angle which this joining line makes with the axis of x we will call ϕ . If y is the ordinate of the point on the hyperbola, then evidently

$$y = \frac{b^2 \tan \phi}{\sqrt{a^2 + b^2}},$$

and

$$x = \frac{a}{\cos \phi} \sqrt{1 - \frac{a^2 \sin^2 \phi}{a^2 + b^2}} = \frac{a}{\cos \phi} \sqrt{1 - \frac{1}{e^2} \sin^2 \phi};$$

whence

$$\begin{aligned} s &= \int \sqrt{ax^2 + dy^2} = \frac{b^2}{c} \int_0^\phi \frac{d\phi}{\cos^2 \phi \sqrt{1 - \frac{1}{e^2} \sin^2 \phi}} \\ &= \frac{b^2}{c} \int_0^\phi \frac{d\phi}{\cos^2 \phi \sqrt{1 - k^2 \sin^2 \phi}}. \end{aligned}$$

But

$$\begin{aligned} d(\tan \phi \sqrt{1 - k^2 \sin^2 \phi}) &= d\phi \sqrt{1 - k^2 \sin^2 \phi} + d\phi \frac{1 - k^2}{\sqrt{1 - e^2 \sin^2 \phi}} \\ &\quad - \frac{1 - k^2}{\cos^2 \phi \sqrt{1 - e^2 \sin^2 \phi}} d\phi. \end{aligned}$$

Consequently

$$\begin{aligned} s &= \frac{b^2}{c} \int_0^\phi \frac{d\phi}{\cos^2 \phi \sqrt{1 - k^2 \sin^2 \phi}} \\ &= \frac{b^2}{c} F(k, \phi) - cE(k, \phi) + c \tan \phi \Delta(k, \phi) \\ &= \frac{b^2}{ae} F\left(\frac{1}{e}, \phi\right) - aeE\left(\frac{1}{e}, \phi\right) + ae \tan \phi \Delta\left(\frac{1}{e}, \phi\right). \end{aligned}$$

EXAMPLE. Find the length of the arc of the hyperbola $\frac{x^2}{20.25} - \frac{y^2}{400} = 1$ from the vertex to the point whose ordinate is $\frac{40}{2.05} \tan 15^\circ$. *Ans.* 5.231184.

EXAMPLE. Find the length of the arc of the hyperbola $\frac{x^2}{144} - \frac{y^2}{81} = 100$ from the vertex to the point whose ordinate is 0.6. *Ans.* 0.6582.

SHORT-TITLE CATALOGUE

OF THE
PUBLICATIONS
OF
JOHN WILEY & SONS,
NEW YORK.

LONDON: CHAPMAN & HALL, LIMITED.

ARRANGED UNDER SUBJECTS.

Descriptive circulars sent on application. Books marked with an asterisk (*) are sold at *net* prices only. All books are bound in cloth unless otherwise stated.

AGRICULTURE.

Armsby's Manual of Cattle-feeding.	12mo, \$1 75
Principles of Animal Nutrition.	8vo, 4 00
Budd and Hansen's American Horticultural Manual:	
Part I. Propagation, Culture, and Improvement.	12mo, 1 50
Part II. Systematic Pomology.	12mo, 1 50
Downing's Fruits and Fruit-trees of America.	8vo, 5 00
Elliott's Engineering for Land Drainage.	12mo, 1 50
Practical Farm Drainage.	12mo, 1 00
Graves's Forest Mensuration.	8vo, 4 00
Green's Principles of American Forestry.	12mo, 1 50
Grotenfelt's Principles of Modern Dairy Practice. (Woll.).	12mo, 2 00
Kemp's Landscape Gardening.	12mo, 2 50
Maynard's Landscape Gardening as Applied to Home Decoration.	12mo, 1 50
* McKay and Larsen's Principles and Practice of Butter-making.	8vo, 1 50
Sanderson's Insects Injurious to Staple Crops.	12mo, 1 50
Insects Injurious to Garden Crops. (In preparation.)	
Insects Injuring Fruits. (In preparation.)	
Stockbridge's Rocks and Soils.	8vo, 2 50
Winton's Microscopy of Vegetable Foods.	8vo, 7 50
Woll's Handbook for Farmers and Dairymen.	16mo, 1 50

ARCHITECTURE.

Baldwin's Steam Heating for Buildings.	12mo, 2 50
Bashore's Sanitation of a Country House.	12mo, 1 00
Berg's Buildings and Structures of American Railroads.	4to, 5 00
Birkmire's Planning and Construction of American Theatres.	8vo, 3 00
Architectural Iron and Steel.	8vo, 3 50
Compound Riveted Girders as Applied in Buildings.	8vo, 2 00
Planning and Construction of High Office Buildings.	8vo, 3 50
Skeleton Construction in Buildings.	8vo, 3 00
Brigg's Modern American School Buildings.	8vo, 4 00

Carpenter's Heating and Ventilating of Buildings.	.8vo,	4 00
Freitag's Architectural Engineering.	.8vo,	3 50
Fireproofing of Steel Buildings.	.8vo,	2 50
French and Ives's Stereotomy.	.8vo,	2 50
Gerhard's Guide to Sanitary House-inspection.	.16mo,	1 00
Theatre Fires and Panics.	.12mo,	1 50
*Greene's Structural Mechanics.	.8vo,	2 50
Holly's Carpenters' and Joiners' Handbook.	.18mo,	75
Johnson's Statics by Algebraic and Graphic Methods.	.8vo,	2 00
Kidder's Architects' and Builders' Pocket-book. Rewritten Edition.	.16mo, mor.,	5 00
Merrill's Stones for Building and Decoration.	.8vo,	5 00
Non-metallic Minerals: Their Occurrence and Uses.	.8vo,	4 00
Monckton's Stair-building.	.4to,	4 00
Patton's Practical Treatise on Foundations.	.8vo,	5 00
Peabody's Naval Architecture.	.8vo,	7 50
Rice's Concrete-block Manufacture.	.8vo,	2 00
Richey's Handbook for Superintendents of Construction.	.16mo, mor.,	4 00
* Building Mechanics' Ready Reference Book. Carpenters' and Wood- workers' Edition.	.16mo, morocco,	1 50
Sabin's Industrial and Artistic Technology of Paints and Varnish.	.8vo,	3 00
Siebert and Biggin's Modern Stone-cutting and Masonry.	.8vo,	1 50
Snow's Principal Species of Wood.	.8vo,	3 50
Sondericker's Graphic Statics with Applications to Trusses, Beams, and Arches. 8vo,	.8vo,	2 00
Towne's Locks and Builders' Hardware.	.18mo, morocco,	3 00
Wait's Engineering and Architectural Jurisprudence.	.8vo,	6 00
Law of Operations Preliminary to Construction in Engineering and Archi- tecture.	.8vo,	5 00
Law of Contracts.	.8vo,	3 00
Wood's Rustless Coatings: Corrosion and Electrolysis of Iron and Steel.	.8vo,	4 00
Worcester and Atkinson's Small Hospitals, Establishment and Maintenance, Suggestions for Hospital Architecture, with Plans for a Small Hospital. 12mo,	.12mo,	1 25
The World's Columbian Exposition of 1893.	Large 4to,	1 00

ARMY AND NAVY.

Bernadou's Smokeless Powder, Nitro-cellulose, and the Theory of the Cellulose Molecule.	.12mo,	2 50
* Bruff's Text-book Ordnance and Gunnery.	.8vo,	6 00
Chase's Screw Propellers and Marine Propulsion.	.8vo,	3 00
Cloke's Gunner's Examiner.	.8vo,	1 50
Craig's Azimuth.	.4to,	3 50
Crehore and Squier's Polarizing Photo-chronograph.	.8vo,	3 00
* Davis's Elements of Law.	.8vo,	2 50
* Treatise on the Military Law of United States.	.8vo,	7 00
De Brack's Cavalry Outposts Duties. (Carr.).	.24mo, morocco,	2 00
Dietz's Soldier's First Aid Handbook.	.16mo, morocco,	1 25
* Dudley's Military Law and the Procedure of Courts-martial.	Large 12mo,	2 50
Durand's Resistance and Propulsion of Ships.	.8vo,	5 00
* Dyer's Handbook of Light Artillery.	.12mo,	3 00
Eissler's Modern High Explosives.	.8vo,	4 00
* Fieberger's Text-book on Field Fortification.	Small 8vo,	2 00
Hamilton's The Gunner's Catechism.	.18mo,	1 00
* Hoff's Elementary Naval Tactics.	.8vo,	1 50

Ingalls's Handbook of Problems in Direct Fire.	8vo,	4 00
* Ballistic Tables.	8vo,	1 50
* Lyons's Treatise on Electromagnetic Phenomena. Vols. I. and II. 8vo, each,	8vo, each,	6 00
* Mahan's Permanent Fortifications. (Mercur.)	8vo, half morocco,	7 50
Manual for Courts-martial.	16mo, morocco,	1 50
* Mercur's Attack of Fortified Places.	12mo,	2 00
* Elements of the Art of War.	8vo,	4 00
Metcalf's Cost of Manufactures—And the Administration of Workshops.	8vo,	5 00
* Ordnance and Gunnery. 2 vols.	12mo,	5 00
Murray's Infantry Drill Regulations.	18mo, paper,	10
Nixon's Adjutants' Manual.	24mo,	1 00
Peabody's Naval Architecture.	8vo,	7 50
* Phelps's Practical Marine Surveying.	8vo,	2 50
Powell's Army Officer's Examiner.	12mo,	4 00
Sharpe's Art of Subsisting Armies in War.	18mo, morocco,	1 50
* Tupes and Poole's Manual of Bayonet Exercises and Musketry Fencing.	24mo, leather,	50
* Walke's Lectures on Explosives.	8vo,	4 00
Weaver's Military Explosives.	8vo,	3 00
* Wheeler's Siege Operations and Military Mining.	8vo,	2 00
Winthrop's Abridgment of Military Law.	12mo,	2 50
Woodhull's Notes on Military Hygiene.	16mo,	1 50
Young's Simple Elements of Navigation.	16mo, morocco,	2 00

ASSAYING.

Fletcher's Practical Instructions in Quantitative Assaying with the Blowpipe.	12mo, morocco,	1 50
Furman's Manual of Practical Assaying.	8vo,	3 00
Lodge's Notes on Assaying and Metallurgical Laboratory Experiments.	8vo,	3 00
Low's Technical Methods of Ore Analysis.	8vo,	3 00
Miller's Manual of Assaying.	12mo,	1 00
Cyanide Process.	12mo,	1 00
Minet's Production of Aluminum and its Industrial Use. (Waldo.)	12mo,	2 50
O'Driscoll's Notes on the Treatment of Gold Ores.	8vo,	2 00
Ricketts and Miller's Notes on Assaying.	8vo,	3 00
Robinet and Lenglen's Cyanide Industry. (Le Clerc.)	8vo,	4 00
Ulke's Modern Electrolytic Copper Refining.	8vo,	3 00
Wilson's Cyanide Processes.	12mo,	1 50
Chlorination Process.	12mo,	1 50

ASTRONOMY.

Comstock's Field Astronomy for Engineers.	8vo,	2 50
Craig's Azimuth.	4to,	3 50
Crandall's Text-book on Geodesy and Least Squares.	8vo,	3 00
Doolittle's Treatise on Practical Astronomy.	8vo,	4 00
Gore's Elements of Geodesy.	8vo,	2 50
Hayford's Text-book of Geodetic Astronomy.	8vo,	3 00
Merriman's Elements of Precise Surveying and Geodesy.	8vo,	2 50
* Michie and Harlow's Practical Astronomy.	8vo,	3 00
* White's Elements of Theoretical and Descriptive Astronomy	12mo	00

BOTANY.

Davenport's Statistical Methods, with Special Reference to Biological Variation.	16mo, morocco,	1 25
Thomé and Bennett's Structural and Physiological Botany.	16mo,	2 25
Westermaier's Compendium of General Botany. (Schneider.)	8vo,	2 00

CHEMISTRY.

* Abegg's Theory of Electrolytic Dissociation. (Von Ende.)	12mo,	1 25
Adriance's Laboratory Calculations and Specific Gravity Tables.	12mo,	1 25
Alexeyeff's General Principles of Organic Synthesis. (Matthews.)	8vo,	3 00
Allen's Tables for Iron Analysis.	8vo,	3 00
Arnold's Compendium of Chemistry. (Mandel.)	Small 8vo,	3 50
Austen's Notes for Chemical Students	12mo,	1 50
Bernadou's Smokeless Powder.—Nitro-cellulose, and Theory of the Cellulose Molecule.	12mo,	2 50
* Browning's Introduction to the Rarer Elements.	8vo,	1 50
Brush and Penfield's Manual of Determinative Mineralogy.	8vo,	4 00
* Claassen's Beet-sugar Manufacture. (Hall and Rolfe.)	8vo,	3 00
Classen's Quantitative Chemical Analysis by Electrolysis. (Eoltwood.)	8vo,	3 00
Cohn's Indicators and Test-papers.	12mo,	2 00
Tests and Reagents.	8vo,	3 00
Crafts's Short Course in Qualitative Chemical Analysis. (Schaeffer.)	12mo,	1 50
* Danneef's Electrochemistry. (Merriam.)	12mo,	1 25
Dolezalek's Theory of the Lead Accumulator (Storage Battery). (Von Ende.)	12mo,	2 50
Drechsel's Chemical Reactions. (Merrill.)	12mo,	1 25
Duhem's Thermodynamics and Chemistry. (Burgess.)	8vo,	4 00
Eissler's Modern High Explosives.	8vo,	4 00
Effront's Enzymes and their Applications. (Prescott.)	8vo,	3 00
Erdmann's Introduction to Chemical Preparations. (Dunlap.)	12mo,	1 25
Fletcher's Practical Instructions in Quantitative Assaying with the Blowpipe.	12mo, morocco,	1 50
Fowler's Sewage Works Analyses.	12mo,	2 00
Fresenius's Manual of Qualitative Chemical Analysis. (Wells.)	8vo,	5 00
Manual of Qualitative Chemical Analysis. Part I. Descriptive. (Wells.)	8vo,	3 00
System of Instruction in Quantitative Chemical Analysis. (Cohn.)	2 vols. 8vo,	12 50
Fuertes's Water and Public Health.	12mo,	1 50
Furman's Manual of Practical Assaying.	8vo,	3 00
* Getman's Exercises in Physical Chemistry.	12mo,	2 00
Gill's Gas and Fuel Analysis for Engineers.	12mo,	1 25
* Gooch and Browning's Outlines of Qualitative Chemical Analysis. Small	8vo,	1 25
Grotenfelt's Principles of Modern Dairy Practice. (Woll.)	12mo,	2 00
Groth's Introduction to Chemical Crystallography (Marshall)	12mo,	1 25
Hammarsten's Text-book of Physiological Chemistry. (Mandel.)	8vo,	4 00
Helm's Principles of Mathematical Chemistry. (Morgan.)	12mo,	1 50
Hering's Ready Reference Tables (Conversion Factors).	16mo, morocco,	2 50
Hind's Inorganic Chemistry.	8vo,	3 00
* Laboratory Manual for Students	12mo,	1 00
Holleman's Text-book of Inorganic Chemistry. (Cooper.)	8vo,	2 50
Text-book of Organic Chemistry. (Walker and Mott.)	8vo,	2 50
* Laboratory Manual of Organic Chemistry. (Walker.)	12mo,	1 00
Hopkins's Oil-chemists' Handbook.	8vo,	3 00
Iddings's Rock Minerals.	8vo,	5 00
Jackson's Directions for Laboratory Work in Physiological Chemistry.	8vo,	1 25
Keep's Cast Iron.	8vo,	2 50
Ladd's Manual of Quantitative Chemical Analysis.	12mo,	1 00
Landauer's Spectrum Analysis. (Tingle.)	8vo,	3 00
* Langworthy and Austen. The Occurrence of Aluminium in Vegetable Products, Animal Products, and Natural Waters.	8vo,	2 00
Lassar-Cohn's Application of Some General Reactions to Investigations in Organic Chemistry. (Tingle.)	12mo,	1 00
Leach's The Inspection and Analysis of Food with Special Reference to State Control.	8vo,	7 50
Löb's Electrochemistry of Organic Compounds. (Lorenz.)	8vo,	3 00

Van Deventer's Physical Chemistry for Beginners. (Boltwood.)	12mo,	1 50
* Walke's Lectures on Explosives.	8vo,	4 00
Ware's Beet-sugar Manufacture and Refining.	Small 8vo, cloth,	4 00
Washington's Manual of the Chemical Analysis of Rocks.	8vo,	2 00
Weaver's Military Explosives.	8vo,	3 00
Wehrenfennig's Analysis and Softening of Boiler Feed-Water	8vo,	4 00
Wells's Laboratory Guide in Qualitative Chemical Analysis.	8vo,	1 50
Short Course in Inorganic Qualitative Chemical Analysis for Engineering Students.	12mo,	1 50
Text-book of Chemical Arithmetic	12mo,	1 25
Whipple's Microscopy of Drinking-water.	8vo,	3 50
Wilson's Cyanide Processes	12mo,	1 50
Chlorination Process.	12mo,	1 50
Winton's Microscopy of Vegetable Foods.	8vo,	7 50
Wulling's Elementary Course in Inorganic, Pharmaceutical, and Medical Chemistry.	12mo,	2 00

CIVIL ENGINEERING.

BRIDGES AND ROOFS. HYDRAULICS. MATERIALS OF ENGINEERING. RAILWAY ENGINEERING.

Baker's Engineers' Surveying Instruments.	12mo,	3 00
Bixby's Graphical Computing Table.	Paper 19½ × 24½ inches.	25
Breed and Hosmer's Principles and Practice of Surveying.	8vo,	3 00
* Burr's Ancient and Modern Engineering and the Isthmian Canal	8vo,	3 50
Comstock's Field Astronomy for Engineers.	8vo,	2 50
Crandall's Text-book on Geodesy and Least Squares	8vo,	3 00
Davis's Elevation and Stadia Tables.	8vo,	1 00
Elliott's Engineering for Land Drainage.	12mo,	1 50
Practical Farm Drainage.	12mo,	1 00
* Fiebigger's Treatise on Civil Engineering.	8vo,	5 00
Flemer's Phototopographic Methods and Instruments.	8vo,	5 00
Folwell's Sewerage. (Designing and Maintenance.)	8vo,	3 00
Freitag's Architectural Engineering. 2d Edition, Rewritten	8vo,	3 50
French and Ives's Stereotomy	8vo,	2 50
Goodhue's Municipal Improvements.	12mo,	1 75
Gore's Elements of Geodesy.	8vo,	2 50
Hayford's Text-book of Geodetic Astronomy.	8vo,	3 00
Hering's Ready Reference Tables (Conversion Factors).	16mo, morocco,	2 50
Howe's Retaining Walls for Earth.	12mo,	1 25
* Ives's Adjustments of the Engineer's Transit and Level.	16mo, Eds.	25
Ives and Hilts's Problems in Surveying	16mo, morocco,	1 50
Johnson's (J. B.) Theory and Practice of Surveying.	Small 8vo,	4 00
Johnson's (L. J.) Statics by Algebraic and Graphic Methods.	8vo,	2 00
Laplace's Philosophical Essay on Probabilities. (Truscott and Emory.)	12mo,	2 00
Mahan's Treatise on Civil Engineering. (1873.) (Wood.)	8vo,	5 00
* Descriptive Geometry.	8vo,	1 50
Merriman's Elements of Precise Surveying and Geodesy.	8vo,	2 50
Merriman and Brooks's Handbook for Surveyors.	16mo, morocco,	2 00
Nugent's Plane Surveying.	8vo,	3 50
Ogden's Sewer Design.	12mo,	2 00
Parsons's Disposal of Municipal Refuse.	8vo,	2 00
Patton's Treatise on Civil Engineering.	8vo half leather,	7 50
Reed's Topographical Drawing and Sketching	4to,	5 00
Rideal's Sewage and the Bacterial Purification of Sewage.	8vo,	4 00
Siebert and Biggin's Modern Stone-cutting and Masonry.	8vo,	1 50

Smith's Manual of Topographical Drawing. (McMillan.)	8vo,	2 50
Sondericker's Graphic Statics, with Applications to Trusses, Beams, and Arches.	8vo,	2 00
Taylor and Thompson's Treatise on Concrete, Plain and Reinforced.	8vo,	5 00
* Trautwine's Civil Engineer's Pocket-book.	16mo, morocco,	5 00
Venable's Garbage Crematories in America.	8vo,	2 00
Wait's Engineering and Architectural Jurisprudence.	8vo	6 00
	Sheep,	6 50
Law of Operations Preliminary to Construction in Engineering and Architecture.	8vo,	5 00
	Sheep,	5 50
Law of Contracts.	8vo,	3 00
Warren's Stereotomy—Problems in Stone-cutting.	8vo,	2 50
Webb's Problems in the Use and Adjustment of Engineering Instruments.	16mo, morocco,	1 25
Wilson's Topographic Surveying.	8vo,	3 50

BRIDGES AND ROOFS.

Boller's Practical Treatise on the Construction of Iron Highway Bridges.	8vo,	2 00
* Thames River Bridge.	4to, paper,	5 00
Burr's Course on the Stresses in Bridges and Roof Trusses, Arched Ribs, and Suspension Bridges.	8vo,	3 50
Burr and Falk's Influence Lines for Bridge and Roof Computations.	8vo,	3 00
Design and Construction of Metallic Bridges.	8vo	5 00
Du Bois's Mechanics of Engineering. Vol. II.	Small 4to,	10 00
Foster's Treatise on Wooden Trestle Bridges.	4to,	5 00
Fowler's Ordinary Foundations.	8vo,	3 50
Greene's Roof Trusses.	8vo,	1 25
Bridge Trusses.	8vo,	2 50
Arches in Wood, Iron, and Stone.	8vo	2 50
Howe's Treatise on Arches.	8vo,	4 00
Design of Simple Roof-trusses in Wood and Steel.	8vo,	2 00
Symmetrical Masonry Arches.	8vo,	2 50
Johnson, Bryan, and Turneaure's Theory and Practice in the Designing of Modern Framed Structures.	Small 4to,	10 00
Merriman and Jacoby's Text-book on Roofs and Bridges:		
Part I. Stresses in Simple Trusses.	8vo,	2 50
Part II. Graphic Statics.	8vo,	2 50
Part III. Bridge Design.	8vo,	2 50
Part IV. Higher Structures.	8vo,	2 50
Morison's Memphis Bridge.	4to,	10 00
Waddell's De Pontibus, a Pocket-book for Bridge Engineers.	16mo, morocco,	2 00
* Specifications for Steel Bridges.	12mo,	50
Wright's Designing of Draw-spans. Two parts in one volume.	8vo,	3 50

HYDRAULICS.

Barnes's Ice Formation.	8vo,	3 00
Bazin's Experiments upon the Contraction of the Liquid Vein Issuing from an Orifice. (Trautwine.)	8vo,	2 00
Bovey's Treatise on Hydraulics.	8vo,	5 00
Church's Mechanics of Engineering.	8vo,	6 00
Diagrams of Mean Velocity of Water in Open Channels.	paper,	1 50
Hydraulic Motors.	8vo,	2 00
Coffin's Graphical Solution of Hydraulic Problems.	16mo, morocco,	2 50
Flather's Dynamometers, and the Measurement of Power.	12mo,	3 00

Folwell's Water-supply Engineering.	8vo,	4 00
Frizell's Water-power.	8vo,	5 00
Fuertes's Water and Public Health.	12mo,	1 50
Water-filtration Works.	12mo,	2 50
Ganguillet and Kutter's General Formula for the Uniform Flow of Water in Rivers and Other Channels. (Hering and Trautwine.)	8vo,	4 00
Hazen's Filtration of Public Water-supply.	8vo,	3 00
Hazlehurst's Towers and Tanks for Water-works.	8vo,	2 50
Herschel's 115 Experiments on the Carrying Capacity of Large, Riveted, Metal Conduits.	8vo,	2 00
Mason's Water-supply. (Considered Principally from a Sanitary Standpoint.)	8vo,	4 00
Merriman's Treatise on Hydraulics.	8vo,	5 00
* Michie's Elements of Analytical Mechanics.	8vo,	4 00
Schuyler's Reservoirs for Irrigation, Water-power, and Domestic Water-supply.	Large 8vo,	5 00
* Thomas and Watt's Improvement of Rivers.	4to,	6 00
Turneure and Russell's Public Water-supplies	8vo,	5 00
Wegmann's Design and Construction of Dams.	4to,	5 00
Water-supply of the City of New York from 1658 to 1895.	4to,	10 00
Whipple's Value of Pure Water.	Large 12mo,	1 00
Williams and Hazen's Hydraulic Tables.	8vo,	1 50
Wilson's Irrigation Engineering.	Small 8vo,	4 00
Wolff's Windmill as a Prime Mover.	8vo,	3 00
Wood's Turbines.	8vo,	2 50
Elements of Analytical Mechanics.	8vo,	3 00

MATERIALS OF ENGINEERING.

Baker's Treatise on Masonry Construction.	8vo,	5 00
Roads and Pavements.	8vo,	5 00
Black's United States Public Works	Oblong 4to,	5 00
* Bovey's Strength of Materials and Theory of Structures.	8vo,	7 50
Burr's Elasticity and Resistance of the Materials of Engineering.	8vo,	7 50
Byrne's Highway Construction.	8vo,	5 00
Inspection of the Materials and Workmanship Employed in Construction.	16mo,	3 00
Church's Mechanics of Engineering.	8vo,	6 00
Du Bois's Mechanics of Engineering. Vol. I.	Small 4to,	7 50
* Eckel's Cements, Limes, and Plasters.	8vo,	6 00
Johnson's Materials of Construction.	Large 8vo,	6 00
Fowler's Ordinary Foundations.	8vo,	3 50
Graves's Forest Mensuration.	8vo,	4 00
* Greene's Structural Mechanics.	8vo,	2 50
Keep's Cast Iron.	8vo,	2 50
Lanza's Applied Mechanics.	8vo,	7 50
Marten's Handbook on Testing Materials. (Henning.) 2 vols.	8vo,	7 50
Maurer's Technical Mechanics.	8vo,	4 00
Merrill's Stones for Building and Decoration.	8vo,	5 00
Merriman's Mechanics of Materials.	8vo,	5 00
* Strength of Materials	12mo,	1 00
Metcalfe's Steel. A Manual for Steel-users.	12mo,	2 00
Patton's Practical Treatise on Foundations.	8vo,	5 00
Richardson's Modern Asphalt Pavements.	8vo,	3 00
Richey's Handbook for Superintendents of Construction.	16mo, mor.,	4 00
* Ries's Clays: Their Occurrence, Properties, and Uses.	8vo,	5 00
Rockwell's Roads and Pavements in France.	12mo,	1 25

Sabin's Industrial and Artistic Technology of Paints and Varnish.....	.8vo,	3 00
Smith's Materials of Machines.....	12mo,	1 00
Snow's Principal Species of Wood.....	.8vo,	3 50
Spalding's Hydraulic Cement.....	12mo,	2 00
Text-book on Roads and Pavements.....	12mo,	2 00
Taylor and Thompson's Treatise on Concrete, Plain and Reinforced.....	.8vo,	5 00
Thurston's Materials of Engineering. 3 Parts.....	.8vo,	8 00
Part I. Non-metallic Materials of Engineering and Metallurgy.....	.8vo,	2 00
Part II. Iron and Steel.....	.8vo,	3 50
Part III. A Treatise on Brasses, Bronzes, and Other Alloys and their Constituents.....	.8vo,	2 50
Tillson's Street Pavements and Paving Materials.....	.8vo,	4 00
Waddell's De Pontibus. (A Pocket-book for Bridge Engineers.)..	16mo, mor.,	2 00
* Specifications for Steel Bridges.....	12mo,	50
Wood's (De V.) Treatise on the Resistance of Materials, and an Appendix on the Preservation of Timber.....	.8vo,	2 00
Wood's (De V.) Elements of Analytical Mechanics.....	.8vo,	3 00
Wood's (M. P.) Rustless Coatings: Corrosion and Electrolysis of Iron and Steel.....	.8vo,	4 00

RAILWAY ENGINEERING.

Andrew's Handbook for Street Railway Engineers.....	315 inches, morocco,	1 25
Berg's Buildings and Structures of American Railroads.....	4to,	5 00
Brook's Handbook of Street Railroad Location.....	16mo, morocco,	1 50
Butt's Civil Engineer's Field-book.....	16mo, morocco,	2 50
Crandall's Transition Curve.....	16mo, morocco,	1 50
Railway and Other Earthwork Tables.....	.8vo,	1 50
Dawson's "Engineering" and Electric Traction Pocket-book.....	16mo, morocco,	5 00
Dredge's History of the Pennsylvania Railroad: (1879).....	Paper,	5 00
Fisher's Table of Cubic Yards.....	Cardboard,	25
Godwin's Railroad Engineers' Field-book and Explorers' Guide.....	16mo, mor.,	2 50
Hudson's Tables for Calculating the Cubic Contents of Excavations and Em- bankments.....	.8vo,	1 00
Molitor and Beard's Manual for Resident Engineers.....	16mo,	1 00
Nagle's Field Manual for Railroad Engineers.....	16mo, morocco,	3 00
Philbrick's Field Manual for Engineers.....	16mo, morocco,	3 00
Searles's Field Engineering.....	16mo, morocco,	3 00
Railroad Spiral.....	16mo, morocco,	1 50
Taylor's Prismoidal Formulæ and Earthwork.....	.8vo,	1 50
* Trautwine's Method of Calculating the Cube Contents of Excavations and Embankments by the Aid of Diagrams.....	.8vo,	2 00
The Field Practice of Laying Out Circular Curves for Railroads.....	12mo, morocco,	2 50
Cross-section Sheet.....	Paper,	25
Webb's Railroad Construction.....	16mo, morocco,	5 00
Economics of Railroad Construction.....	Large 12mo,	2 50
Wellington's Economic Theory of the Location of Railways.....	Small 8vo,	5 00

DRAWING.

Barr's Kinematics of Machinery.....	.8vo,	2 50
* Bartlett's Mechanical Drawing.....	.8vo,	3 00
* " " " " Abridged Ed.....	.8vo,	1 50
Coolidge's Manual of Drawing.....	.8vo, paper,	1 00

Coolidge and Freeman's Elements of General Drafting for Mechanical Engineers.....	Oblong 4to,	2 50
Durley's Kinematics of Machines.....	8vo,	4 00
Emch's Introduction to Projective Geometry and its Applications.....	8vo,	2 50
Hill's Text-book on Shades and Shadows, and Perspective.....	8vo,	2 00
Jamison's Elements of Mechanical Drawing.....	8vo,	2 50
Advanced Mechanical Drawing.....	8vo,	2 00
Jones's Machine Design:		
Part I. Kinematics of Machinery.....	8vo,	1 50
Part II. Form, Strength, and Proportions of Parts.....	8vo,	3 00
MacCord's Elements of Descriptive Geometry.....	8vo,	3 00
Kinematics; or, Practical Mechanism.....	8vo,	5 00
Mechanical Drawing.....	4to,	4 00
Velocity Diagrams.....	8vo,	1 50
MacLeod's Descriptive Geometry.....	Small 8vo,	1 50
* Mahan's Descriptive Geometry and Stone-cutting.....	8vo,	1 50
Industrial Drawing. (Thompson.).....	8vo,	3 50
Moyer's Descriptive Geometry.....	8vo,	2 00
Reed's Topographical Drawing and Sketching.....	4to,	5 00
Reid's Course in Mechanical Drawing.....	8vo,	2 00
Text-book of Mechanical Drawing and Elementary Machine Design.....	8vo,	3 00
Robinson's Principles of Mechanism.....	8vo,	3 00
Schwamb and Merrill's Elements of Mechanism.....	8vo,	3 00
Smith's (R. S.) Manual of Topographical Drawing. (McMillan.).....	8vo,	2 50
Smith (A. W.) and Marx's Machine Design.....	8vo,	3 00
* Titsworth's Elements of Mechanical Drawing.....	Oblong 8vo,	1 25
Warren's Elements of Plane and Solid Free-hand Geometrical Drawing.....	12mo,	1 00
Drafting Instruments and Operations.....	12mo,	1 25
Manual of Elementary Projection Drawing.....	12mo,	1 50
Manual of Elementary Problems in the Linear Perspective of Form and Shadow.....	12mo,	1 00
Plane Problems in Elementary Geometry.....	12mo,	1 25
Primary Geometry.....	12mo,	75
Elements of Descriptive Geometry, Shadows, and Perspective.....	8vo,	3 50
General Problems of Shades and Shadows.....	8vo,	3 00
Elements of Machine Construction and Drawing.....	8vo,	7 50
Problems, Theorems, and Examples in Descriptive Geometry.....	8vo,	2 50
Weisbach's Kinematics and Power of Transmission. (Hermann and Klein.).....	8vo,	5 00
Whelpley's Practical Instruction in the Art of Letter Engraving.....	12mo,	2 00
Wilson's (H. M.) Topographic Surveying.....	8vo,	3 50
Wilson's (V. T.) Free-hand Perspective.....	8vo,	2 50
Wilson's (V. T.) Free-hand Lettering.....	8vo,	1 00
Woolf's Elementary Course in Descriptive Geometry.....	Large 8vo,	3 00

ELECTRICITY AND PHYSICS.

* Abegg's Theory of Electrolytic Dissociation. (Von Ende.).....	12mo,	1 25
Anthony and Brackett's Text-book of Physics. (Magie.).....	Small 8vo	3 00
Anthony's Lecture-notes on the Theory of Electrical Measurements.....	12mo,	1 00
Benjamin's History of Electricity.....	8vo,	3 00
Voltaic Cell.....	8vo,	3 00
Classen's Quantitative Chemical Analysis by Electrolysis. (Boltwood.).....	8vo,	3 00
* Collins's Manual of Wireless Telegraphy.....	12mo,	1 50
Morocco,		2 00
Crehore and Squier's Polarizing Photo-chronograph.....	8vo,	3 00
* Danneel's Electrochemistry. (Merriam.).....	12mo,	1 25
Dawson's "Engineering" and Electric Traction Pocket-book.....	16mo, morocco,	5 00

Dolezalek's Theory of the Lead Accumulator (Storage Battery). (Von Ende.).	12mo,	2 50
Duhem's Thermodynamics and Chemistry. (Burgess.).	8vo,	4 00
Flather's Dynamometers, and the Measurement of Power.	12mo,	3 00
Gilbert's De Magnete. (Mottelay.).	8vo,	2 50
Hanchett's Alternating Currents Explained.	12mo,	1 00
Hering's Ready Reference Tables (Conversion Factors).	16mo morocco,	2 50
Holman's Precision of Measurements.	8vo,	2 00
Telescopic Mirror-scale Method, Adjustments, and Tests.	Large 8vo,	75
Kinzbrunner's Testing of Continuous-current Machines.	8vo,	2 00
Landauer's Spectrum Analysis. (Tingle.).	8vo,	3 00
Le Chatelier's High-temperature Measurements. (Boudouard—Burgess.).	12mo,	3 00
Löb's Electrochemistry of Organic Compounds. (Lorenz.).	8vo,	3 00
* Lyons's Treatise on Electromagnetic Phenomena. Vols. I. and II.	8vo, each,	6 00
* Michie's Elements of Wave Motion Relating to Sound and Light.	8vo,	4 00
Niaudet's Elementary Treatise on Electric Batteries. (Fishback.).	12mo,	2 50
* Parshall and Hobart's Electric Machine Design.	4to, half morocco,	12 50
Reagan's Locomotives: Simple, Compound, and Electric. [New Edition.	Large 12mo,	3 50
* Rosenberg's Electrical Engineering. (Haldane Gee—Kinzbrunner.).	8vo,	2 00
Ryan, Norris, and Hoxie's Electrical Machinery. Vol. I.	8vo,	2 50
Thurston's Stationary Steam-engines.	8vo,	2 50
* Tillman's Elementary Lessons in Heat.	8vo,	1 50
Tory and Pitcher's Manual of Laboratory Physics.	Small 8vo,	1 00
Ulke's Modern Electrolytic Copper Refining.	8vo,	3 00

LAW.

* Davis's Elements of Law.	8vo,	2 50
* Treatise on the Military Law of United States.	8vo,	7 00
*	Sheep,	7 50
* Dudley's Military Law and the Procedure of Courts-martial.	Large 12mo,	2 50
Manual for Courts-martial.	16mo, morocco,	1 50
Wait's Engineering and Architectural Jurisprudence.	8vo,	6 00
	Sheep,	6 50
Law of Operations Preliminary to Construction in Engineering and Architecture.	8vo	5 00
	Sheep,	5 50
Law of Contracts.	8vo,	3 00
Winthrop's Abridgment of Military Law.	12mo,	2 50

MANUFACTURES.

Bernadou's Smokeless Powder—Nitro-cellulose and Theory of the Cellulose Molecule.	12mo,	2 50
Bolland's Iron Founder.	12mo,	2 50
The Iron Founder," Supplement.	12mo,	2 50
Encyclopedia of Founding and Dictionary of Foundry Terms Used in the Practice of Moulding.	12mo,	3 00
* Claassen's Beet-sugar Manufacture. (Hall and Rolfe.).	8vo,	3 00
* Eckel's Cements, Limes, and Plasters.	8vo,	6 00
Eissler's Modern High Explosives.	8vo,	4 00
Effront's Enzymes and their Applications. (Prescott.).	8vo,	3 00
Fitzgerald's Boston Machinist.	12mo,	1 00
Ford's Boiler Making for Boiler Makers.	18mo,	1 00
Hopkin's Oil-chemists' Handbook.	8vo,	3 00
Keep's Cast Iron.	8vo,	2 50

Leach's The Inspection and Analysis of Food with Special Reference to State Control.....	Large 8vo,	7 50
* McKay and Larsen's Principles and Practice of Butter-making.....	8vo,	1 50
Matthews's The Textile Fibres.....	8vo,	3 50
Metcalf's Steel. A Manual for Steel-users.....	12mo,	2 00
Metcalf's Cost of Manufactures—And the Administration of Workshops.....	8vo,	5 00
Meyer's Modern Locomotive Construction.....	4to,	10 00
Morse's Calculations used in Cane-sugar Factories.....	16mo, morocco,	1 50
* Reising's Guide to Piece-dyeing.....	8vo,	25 00
Rice's Concrete-block Manufacture.....	8vo,	2 00
Sabin's Industrial and Artistic Technology of Paints and Varnish.....	8vo,	3 00
Smith's Press-working of Metals.....	8vo,	3 00
Spalding's Hydraulic Cement.....	12mo,	2 00
Spencer's Handbook for Chemists of Beet-sugar Houses.....	16mo morocco,	3 00
Handbook for Cane Sugar Manufacturers.....	16mo morocco,	3 00
Taylor and Thompson's Treatise on Concrete, Plain and Reinforced.....	8vo,	5 00
Thurston's Manual of Steam-boilers, their Designs, Construction and Operation.....	8vo,	5 00
* Walke's Lectures on Explosives.....	8vo,	4 00
Ware's Beet-sugar Manufacture and Refining.....	Small 8vo,	4 00
Weaver's Military Explosives.....	8vo,	3 00
West's American Foundry Practice.....	12mo,	2 50
Moulder's Text-book.....	12mo,	2 50
Wolff's Windmill as a Prime Mover.....	8vo,	3 00
Wood's Rustless Coatings: Corrosion and Electrolysis of Iron and Steel.....	8vo,	4 00

MATHEMATICS.

Baker's Elliptic Functions.....	8vo,	1 50
* Bass's Elements of Differential Calculus.....	12mo,	4 00
Briggs's Elements of Plane Analytic Geometry.....	12mo,	1 00
Compton's Manual of Logarithmic Computations.....	12mo	1 50
Davis's Introduction to the Logic of Algebra.....	8vo,	1 50
* Dickson's College Algebra.....	Large 12mo,	1 50
* Introduction to the Theory of Algebraic Equations.....	Large 12mo,	1 25
Emch's Introduction to Projective Geometry and its Applications.....	8vo	2 50
Halsted's Elements of Geometry.....	8vo,	1 75
Elementary Synthetic Geometry.....	8vo,	1 50
Rational Geometry.....	12mo,	1 75
* Johnson's (J. B.) Three-place Logarithmic Tables: Vest-pocket size, paper,		15
	100 copies for	5 00
* Mounted on heavy cardboard, 8×10 inches,		25
	10 copies for	2 00
Johnson's (W. W.) Elementary Treatise on Differential Calculus.....	Small 8vo,	3 00
Elementary Treatise on the Integral Calculus.....	Small 8vo,	1 50
Johnson's (W. W.) Curve Tracing in Cartesian Co-ordinates.....	12mo,	1 00
Johnson's (W. W.) Treatise on Ordinary and Partial Differential Equations.....		
	Small 8vo,	3 50
Johnson's (W. W.) Theory of Errors and the Method of Least Squares.....	12mo,	1 50
* Johnson's (W. W.) Theoretical Mechanics.....	12mo,	3 00
Laplace's Philosophical Essay on Probabilities. (Truscott and Emory.).....	12mo,	2 00
* Ludlow and Bass. Elements of Trigonometry and Logarithmic and Other Tables.....	8vo,	3 00
Trigonometry and Tables published separately.....	Each,	2 00
* Ludlow's Logarithmic and Trigonometric Tables.....	8vo	1 00
Manning's Irrational Numbers and their Representation by Sequences and Series.....	12mo,	1 25

Mathematical Monographs. Edited by Mansfield Merriman and Robert S. Woodward. Octavo, each			1 00
No. 1. History of Modern Mathematics, by David Eugene Smith.			
No. 2. Synthetic Projective Geometry, by George Bruce Halsted.			
No. 3. Determinants, by Laenas Gifford Weld.			
No. 4. Hyperbolic Functions, by James McMahon.			
No. 5. Harmonic Functions, by William E. Byerly.			
No. 6. Grassmann's Space Analysis, by Edward W. Hyde.			
No. 7. Probability and Theory of Errors, by Robert S. Woodward.			
No. 8. Vector Analysis and Quaternions, by Alexander Macfarlane.			
No. 9. Differential Equations, by William Woolsey Johnson.			
No. 10. The Solution of Equations, by Mansfield Merriman.			
No. 11. Functions of a Complex Variable, by Thomas S. Fiske.			
Maurer's Technical Mechanics.	8vo,	4 00	
Merriman's Method of Least Squares.	8vo,	2 00	
Rice and Johnson's Elementary Treatise on the Differential Calculus. Sm.	8vo,	3 00	
Differential and Integral Calculus. 2 vols. in one.	Small 8vo,	2 50	
* Veblen and Lennes's Introduction to the Real Infinitesimal Analysis of One Variable	8vo,	2 00	
Wood's Elements of Co-ordinate Geometry.	8vo,	2 00	
Trigonometry: Analytical, Plane, and Spherical	12mo,	1 00	

MECHANICAL ENGINEERING.

MATERIALS OF ENGINEERING, STEAM-ENGINES AND BOILERS.

Bacon's Forge Practice.	12mo,	1 50
Baldwin's Steam Heating for Buildings.	12mo,	2 50
Barr's Kinematics of Machinery.	8vo,	2 50
* Bartlett's Mechanical Drawing.	8vo,	3 00
* " " " " Abridged Ed.	8vo,	1 50
Benjamin's Wrinkles and Recipes.	12mo,	2 00
Carpenter's Experimental Engineering.	8vo,	6 00
Heating and Ventilating Buildings.	8vo,	4 00
Clerk's Gas and Oil Engine.	Small 8vo,	4 00
Coolidge's Manual of Drawing.	8vo, paper,	1 00
Coolidge and Freeman's Elements of General Drafting for Mechanical Engineers.	Oblong 4to,	2 50
Cromwell's Treatise on Toothed Gearing.	12mo,	1 50
Treatise on Belts and Pulleys.	12mo,	1 50
Durley's Kinematics of Machines.	8vo,	4 00
Flather's Dynamometers and the Measurement of Power.	12mo,	3 00
Rope Driving.	12mo,	2 00
Gill's Gas and Fuel Analysis for Engineers.	12mo,	1 25
Hall's Car Lubrication.	12mo,	1 00
Hering's Ready Reference Tables (Conversion Factors).	16mo, morocco,	2 50
Hutton's The Gas Engine.	8vo,	5 00
Jamison's Mechanical Drawing.	8vo,	2 50
Jones's Machine Design:		
Part I. Kinematics of Machinery.	8vo,	1 50
Part II. Form, Strength, and Proportions of Parts.	8vo,	3 00
Kent's Mechanical Engineers' Pocket-book.	16mo, morocco,	5 00
Kerr's Power and Power Transmission.	8vo,	2 00
Leonard's Machine Shop, Tools, and Methods.	8vo,	4 00
* Lorenz's Modern Refrigerating Machinery. (Pope, Haven, and Dean.)	8vo,	4 00
MacCord's Kinematics; or, Practical Mechanism.	8vo,	5 00
Mechanical Drawing.	4to,	4 00
Velocity Diagrams.	8vo,	1 50

MacFarland's Standard Reduction Factors for Gases.	8vo,	1 50
Mahan's Industrial Drawing. (Thompson.).....	8vo	3 50
Poole's Caloric Power of Fuels.	8vo,	3 00
Reid's Course in Mechanical Drawing.	8vo,	2 00
Text-book of Mechanical Drawing and Elementary Machine Design.	8vo,	3 00
Richard's Compressed Air.	12mo,	1 50
Robinson's Principles of Mechanism.	8vo,	3 00
Schwarz and Merrill's Elements of Mechanism.	8vo,	3 00
Smith's (O.) Press-working of Metals.	8vo	3 00
Smith (A. W.) and Marx's Machine Design.	8vo,	3 00
Thurston's Treatise on Friction and Lost Work in Machinery and Mill Work.	8vo,	3 00
Animal as a Machine and Prime Motor, and the Laws of Energetics.	12mo,	1 00
Tillson's Complete Automobile Instructor.	16mo,	1 50
	Morocco,	2 00
Warren's Elements of Machine Construction and Drawing.	8vo,	7 50
Weisbach's Kinematics and the Power of Transmission. (Herrmann—Klein.)	8vo,	5 00
Machinery of Transmission and Governors. (Herrmann—Klein.)	8vo,	5 00
Wolf's Windmill as a Prime Mover.	8vo,	3 00
Wood's Turbines.	8vo,	2 50

MATERIALS OF ENGINEERING.

* Bovey's Strength of Materials and Theory of Structures.	8vo,	7 50
Burr's Elasticity and Resistance of the Materials of Engineering. 6th Edition. Reset.	8vo,	7 50
Church's Mechanics of Engineering.	8vo,	6 00
* Greene's Structural Mechanics.	8vo,	2 50
Johnson's Materials of Construction.	8vo,	6 00
Keep's Cast Iron.	8vo,	2 50
Lanza's Applied Mechanics.	8vo,	7 50
Martens's Handbook on Testing Materials. (Henning.)	8vo,	7 50
Maurer's Technical Mechanics.	8vo,	4 00
Merriman's Mechanics of Materials.	8vo,	5 00
* Strength of Materials.	12mo,	1 00
Metcalf's Steel. A Manual for Steel-users.	12mo,	2 00
Sabin's Industrial and Artistic Technology of Paints and Varnish.	8vo,	3 00
Smith's Materials of Machines.	12mo,	1 00
Thurston's Materials of Engineering.	3 vols., 8vo,	8 00
Part II. Iron and Steel.	8vo,	3 50
Part III. A Treatise on Brasses, Bronzes, and Other Alloys and their Constituents.	8vo,	2 50
Wood's (De V.) Treatise on the Resistance of Materials and an Appendix on the Preservation of Timber.	8vo,	2 00
Elements of Analytical Mechanics.	8vo,	3 00
Wood's (M. P.) Rustless Coatings: Corrosion and Electrolysis of Iron and Steel.	8vo,	4 00

STEAM-ENGINES AND BOILERS.

Berry's Temperature-entropy Diagram.	12mo,	1 25
Carnot's Reflections on the Motive Power of Heat. (Thurston.)	12mo,	1 50
Dawson's "Engineering" and Electric Traction Pocket-book.	16mo, mor.,	5 00
Ford's Boiler Making for Boiler Makers.	18mo,	1 00
Goss's Locomotive Sparks.	8vo,	2 00
Locomotive Performance.	8vo,	5 00
Hemenway's Indicator Practice and Steam-engine Economy.	12mo,	2 00

Hutton's Mechanical Engineering of Power Plants.	.8vo,	5 00
Heat and Heat-engines.	.8vo,	5 00
Kent's Steam boiler Economy.	.8vo,	4 00
Kneass's Practice and Theory of the Injector.	.8vo,	1 50
MacCord's Slide-valves.	.8vo,	2 00
Meyer's Modern Locomotive Construction.	.4to,	10 00
Peabody's Manual of the Steam-engine Indicator.	.12mo,	1 50
Tables of the Properties of Saturated Steam and Other Vapors	.8vo,	1 00
Thermodynamics of the Steam-engine and Other Heat-engines.	.8vo,	5 00
Valve-gears for Steam-engines.	.8vo,	2 50
Peabody and Miller's Steam-boilers.	.8vo,	4 00
Pray's Twenty Years with the Indicator.	Large 8vo,	2 50
Pupin's Thermodynamics of Reversible Cycles in Gases and Saturated Vapors. (Osterberg.)	.12mo,	1 25
Reagan's Locomotives: Simple, Compound, and Electric. New Edition. Large 12mo, 3 50		
Rontgen's Principles of Thermodynamics. (Du Bois.)	.8vo,	5 00
Sinclair's Locomotive Engine Running and Management.	.12mo,	2 00
Smart's Handbook of Engineering Laboratory Practice.	.12mo,	2 50
Snow's Steam-boiler Practice.	.8vo,	3 00
Spangler's Valve-gears.	.8vo,	2 50
Notes on Thermodynamics.	.12mo,	1 00
Spangler, Greene, and Marshall's Elements of Steam-engineing	.8vo,	3 00
Thomas's Steam-turbines.	.8vo,	3 50
Thurston's Handy Tables.	.8vo,	1 50
Manual of the Steam-engine.	2 vols., 8vo,	10 00
Part I. History, Structure, and Theory.	.8vo,	6 00
Part II. Design, Construction, and Operation.	.8vo,	6 00
Handbook of Engine and Boiler Trials, and the Use of the Indicator and the Prony Brake.	.8vo,	5 00
Stationary Steam-engines.	.8vo,	2 50
Steam-boiler Explosions in Theory and in Practice	.12mo,	1 50
Manual of Steam-boilers, their Designs, Construction, and Operation. 8vo,		5 00
Wehrenfenning's Analysis and Softening of Boiler Feed-water (Patterson) 8vo,		4 00
Weisbach's Heat, Steam, and Steam-engines. (Du Bois.)	.8vo,	5 00
Whitham's Steam-engine Design.	.8vo,	5 00
Wood's Thermodynamics, Heat Motors, and Refrigerating Machines. .8vo,		4 00

MECHANICS AND MACHINERY.

Barr's Kinematics of Machinery.	.8vo,	2 50
* Bovey's Strength of Materials and Theory of Structures	.8vo,	7 50
Chase's The Art of Pattern-making.	.12mo,	2 50
Church's Mechanics of Engineering.	.8vo,	6 00
Notes and Examples in Mechanics.	.8vo,	2 00
Compton's First Lessons in Metal-working.	.12mo,	1 50
Compton and De Groodt's The Speed Lathe.	.12mo,	1 50
Cromwell's Treatise on Toothed Gearing.	.12mo,	1 50
Treatise on Belts and Pulleys.	.12mo,	1 50
Dana's Text-book of Elementary Mechanics for Colleges and Schools. 12mo,		1 50
Dingey's Machinery Pattern Making	.12mo,	2 00
Dredge's Record of the Transportation Exhibits Building of the World's Columbian Exposition of 1893.	.4to half morocco,	5 00
Du Bois's Elementary Principles of Mechanics:		
Vol. I. Kinematics.	.8vo,	3 50
Vol. II. Statics.	.8vo,	4 00
Mechanics of Engineering. Vol. I.	Small 4to,	7 50
Vol. II.	Small 4to,	10 00
Durley's Kinematics of Machines.	.8vo,	4 00

Fitzgerald's Boston Machinist.	16mo,	1 00
Flather's Dynamometers, and the Measurement of Power.	12mo,	3 00
Rope Driving.	12mo,	2 00
Goss's Locomotive Sparks.	8vo,	2 00
Locomotive Performance.	8vo,	5 00
* Greene's Structural Mechanics.	8vo,	2 50
Hall's Car Lubrication.	12mo,	1 00
Holly's Art of Saw Filing.	18mo,	75
James's Kinematics of a Point and the Rational Mechanics of a Particle.	Small 8vo,	2 00
* Johnson's (W. W.) Theoretical Mechanics.	12mo,	3 00
Johnson's (L. J.) Statics by Graphic and Algebraic Methods.	8vo,	2 00
Jones's Machine Design:		
Part I. Kinematics of Machinery.	8vo,	1 50
Part II. Form, Strength, and Proportions of Parts.	8vo,	3 00
Kerr's Power and Power Transmission.	8vo,	2 00
Lanza's Applied Mechanics.	8vo,	7 50
Leonard's Machine Shop, Tools, and Methods.	8vo,	4 00
* Lorenz's Modern Refrigerating Machinery. (Pope, Haven, and Dean.)	8vo,	4 00
MacCord's Kinematics; or, Practical Mechanism.	8vo,	5 00
Velocity Diagrams.	8vo,	1 50
* Martin's Text Book on Mechanics, Vol. I, Statics.	12mo,	1 25
Maurer's Technical Mechanics.	8vo,	4 00
Merriman's Mechanics of Materials.	8vo,	5 00
* Elements of Mechanics.	12mo,	1 00
* Michie's Elements of Analytical Mechanics.	8vo,	4 00
* Parshall and Hobart's Electric Machine Design.	4to, half morocco,	12 50
Reagan's Locomotives: Simple, Compound, and Electric. New Edition.		
Large 12mo,	3 00	
Reid's Course in Mechanical Drawing.	8vo,	2 00
Text-book of Mechanical Drawing and Elementary Machine Design.	8vo,	3 00
Richards's Compressed Air.	12mo,	1 50
Robinson's Principles of Mechanism.	8vo,	3 00
Ryan, Norris, and Hoxie's Electrical Machinery. Vol. I.	8vo,	2 50
Sanborn's Mechanics: Problems.	Large 12mo,	1 50
Schwamb and Merrill's Elements of Mechanism.	8vo,	3 00
Sinclair's Locomotive-engine Running and Management.	12mo,	2 00
Smith's (O.) Press-working of Metals.	8vo,	3 00
Smith's (A. W.) Materials of Machines.	12mo,	1 00
Smith (A. W.) and Marx's Machine Design.	8vo,	3 00
Spangler, Greene, and Marshall's Elements of Steam-engineering.	8vo,	3 00
Thurston's Treatise on Friction and Lost Work in Machinery and Mill Work.	8vo,	3 00
Animal as a Machine and Prime Motor, and the Laws of Energetics.	12mo,	1 00
Tillson's Complete Automobile Instructor.	16mo,	1 50
Morocco,	2 00	
Warren's Elements of Machine Construction and Drawing.	8vo,	7 50
Weisbach's Kinematics and Power of Transmission. (Herrmann—Klein.)	8vo,	5 00
Machinery of Transmission and Governors. (Herrmann—Klein.)	8vo,	5 00
Wood's Elements of Analytical Mechanics.	8vo,	3 00
Principles of Elementary Mechanics.	12mo,	1 25
Turbines.	8vo,	2 50
The World's Columbian Exposition of 1893.	4to,	1 00

MEDICAL.

De Fursac's Manual of Psychiatry. (Rosanoff and Collins.)....	Large 12mo,	2 50
Ehrlich's Collected Studies on Immunity. (Bolduan.)	8vo,	6 00
Hammarsten's Text-book on Physiological Chemistry. (Mandel.)	8vo,	4 00

Lassar-Cohn's Practical Urinary Analysis. (Lorenz.)	12mo,	1 00
* Pauli's Physical Chemistry in the Service of Medicine. (Fischer.)	12mo,	1 25
* Pozzi-Escot's The Toxins and Venoms and their Antibodies. (Cohn.)	12mo,	1 00
Rostoski's Serum Diagnosis. (Bolduan.)	12mo,	1 00
Salkowski's Physiological and Pathological Chemistry. (Orndorff.)	8vo,	2 50
* Satterlee's Outlines of Human Embryology	12mo,	1 25
Steel's Treatise on the Diseases of the Dog.	8vo,	3 50
Von Behring's Suppression of Tuberculosis. (Bolduan.)	12mo,	1 00
Wassermann's Immune Sera: Hæmolysis, Cytotoxins, and Precipitins. (Bolduan.)	12mo, cloth,	1 00
Woodhull's Notes on Military Hygiene	16mo,	1 50
* Personal Hygiene.	12mo,	1 00
Wulling's An Elementary Course in Inorganic Pharmaceutical and Medical Chemistry	12mo,	2 00

METALLURGY.

Egleston's Metallurgy of Silver, Gold, and Mercury:

Vol. I. Silver.	8vo,	7 50
Vol. II. Gold and Mercury.	8vo,	7 50
Goessel's Minerals and Metals: A Reference Book.	16mo, mor.	3 00
* Iles's Lead-smelting.	12mo,	2 50
Keep's Cast Iron.	8vo,	2 50
Kunhardt's Practice of Ore Dressing in Europe.	8vo,	1 50
Le Chatelier's High-temperature Measurements. (Boudouard—Burgess.)	12mo,	3 00
Metcalf's Steel. A Manual for Steel-users.	12mo,	2 00
Miller's Cyanide Process.	12mo,	1 00
Minet's Production of Aluminum and its Industrial Use. (Waldo.)	12mo,	2 50
Robine and Lenglen's Cyanide Industry. (Le Clerc.)	8vo,	4 00
Smith's Materials of Machines.	12mo,	1 00
Thurston's Materials of Engineering. In Three Parts.	8vo,	8 00
Part II. Iron and Steel.	8vo,	3 50
Part III. A Treatise on Brasses, Bronzes, and Other Alloys and their Constituents.	8vo,	2 50
Ulke's Modern Electrolytic Copper Refining.	8vo,	3 00

MINERALOGY.

Barringer's Description of Minerals of Commercial Value. Oblong, morocco,	2 50
Boyd's Resources of Southwest Virginia.	8vo, 3 00
Map of Southwest Virginia.	Pocket-book form. 2 00
Brush's Manual of Determinative Mineralogy. (Penfield.)	8vo, 4 00
Chester's Catalogue of Minerals.	8vo, paper, 1 00
	Cloth, 1 25
Dictionary of the Names of Minerals.	8vo, 3 50
Dana's System of Mineralogy.	Large 8vo, half leather, 12 50
First Appendix to Dana's New "System of Mineralogy."	Large 8vo, 1 00
Text-book of Mineralogy.	8vo, 4 00
Minerals and How to Study Them	12mo, 1 50
Catalogue of American Localities of Minerals.	Large 8vo, 1 00
Manual of Mineralogy and Petrography.	12mo 2 00
Douglas's Untechnical Addresses on Technical Subjects.	12mo, 1 00
Eakle's Mineral Tables.	8vo, 1 25
Egleston's Catalogue of Minerals and Synonyms.	8vo, 2 50
Goessel's Minerals and Metals: A Reference Book.	16mo, mor. 3 00
Groth's Introduction to Chemical Crystallography (Marshall)	12mo, 1 25

Iddings's Rock Minerals.	8vo,	5 00
Merrill's Non-metallic Minerals: Their Occurrence and Uses.	8vo,	4 00
* Penfield's Notes on Determinative Mineralogy and Record of Mineral Tests.	8vo, paper,	50
* Richards's Synopsis of Mineral Characters.	12mo, morocco,	1 25
* Ries's Clays: Their Occurrence, Properties, and Uses.	8vo,	5 00
Rosenbusch's Microscopical Physiography of the Rock-making Minerals. (Iddings.).	8vo,	5 00
* Tillman's Text-book of Important Minerals and Rocks.	8vo,	2 00

MINING.

Boyd's Resources of Southwest Virginia.	8vo,	3 00
Map of Southwest Virginia.	Pocket-book form	2 00
Douglas's Untechnical Addresses on Technical Subjects.	12mo,	1 00
Eissler's Modern High Explosives.	8vo,	4 00
Goesel's Minerals and Metals: A Reference Book.	16mo, mor.	3 00
Goodyear's Coal-mines of the Western Coast of the United States.	12mo,	2 50
Ihlseng's Manual of Mining.	8vo,	5 00
* Iles's Lead-smelting.	12mo,	2 50
Kunhardt's Practice of Ore Dressing in Europe.	8vo,	1 50
Miller's Cyanide Process.	12mo,	1 00
O'Driscoll's Notes on the Treatment of Gold Ores.	8vo,	2 00
Robine and Lenglen's Cyanide Industry. (Le Clerc.).	8vo,	4 00
* Walke's Lectures on Explosives.	8vo,	4 00
Weaver's Military Explosives.	8vo,	3 00
Wilson's Cyanide Processes.	12mo,	1 50
Chlorination Process.	12mo,	1 50
Hydraulic and Placer Mining.	12mo,	2 00
Treatise on Practical and Theoretical Mine Ventilation.	12mo,	1 25

SANITARY SCIENCE.

Bashore's Sanitation of a Country House.	12mo,	1 00
* Outlines of Practical Sanitation.	12mo,	1 25
Folwell's Sewerage. (Designing, Construction, and Maintenance.).	8vo,	3 00
Water-supply Engineering.	8vo,	4 00
Fowler's Sewage Works Analyses.	12mo,	2 00
Fuertes's Water and Public Health.	12mo,	1 50
Water-filtration Works.	12mo,	2 50
Gerhard's Guide to Sanitary House-inspection.	16mo,	1 00
Hazen's Filtration of Public Water-supplies.	8vo,	3 00
Leach's The Inspection and Analysis of Food with Special Reference to State Control.	8vo,	7 50
Mason's Water-supply. (Considered principally from a Sanitary Standpoint) 8vo,		4 00
Examination of Water. (Chemical and Bacteriological).	12mo,	1 25
* Merriman's Elements of Sanitary Engineering.	8vo,	2 00
Ogden's Sewer Design.	12mo,	2 00
Prescott and Winslow's Elements of Water Bacteriology, with Special Refer- ence to Sanitary Water Analysis.	12mo,	1 25
* Price's Handbook on Sanitation.	12mo,	1 50
Richards's Cost of Food. A Study in Dietsaries.	12mo,	1 00
Cost of Living as Modified by Sanitary Science.	12mo,	1 00
Cost of Shelter.	12mo,	1 00

Richards and Woodman's Air, Water, and Food from a Sanitary Stand-point.	8vo,	2 00
* Richards and Williams's The Dietary Computer.	8vo	1 50
Rideal's Sewage and Bacterial Purification of Sewage.	8vo,	4 00
Turneure and Russell's Public Water-supplies.	8vo,	5 00
Von Behring's Suppression of Tuberculosis. (Bolduan.)	12mo,	1 00
Whipple's Microscopy of Drinking-water.	8vo,	3 50
Winton's Microscopy of Vegetable Foods.	8vo,	7 50
Woodhull's Notes on Military Hygiene.	16mo,	1 50
* Personal Hygiene.	12mo,	1 00

MISCELLANEOUS.

Emmons's Geological Guide-book of the Rocky Mountain Excursion of the International Congress of Geologists.	Large 8vo,	1 5
Ferrel's Popular Treatise on the Winds.	8vo,	4 00
Gannett's Statistical Abstract of the World.	24mo,	75
Haines's American Railway Management.	12mo,	2 50
Ricketts's History of Rensselaer Polytechnic Institute 1824-1894. Small	8vo,	3 00
Rotherham's Emphasized New Testament.	Large 8vo,	2 00
The World's Columbian Exposition of 1893.	4to,	1 00
Winslow's Elements of Applied Microscopy.	12mo,	1 50

HEBREW AND CHALDEE TEXT-BOOKS.

Green's Elementary Hebrew Grammar.	12mo,	1 25
Hebrew Chrestomathy.	8vo,	2 00
Gesenius's Hebrew and Chaldee Lexicon to the Old Testament Scriptures. (Tregelles.).	Small 4to, half morocco,	5 00
Letteris's Hebrew Bible.	8vo,	2 25

QA 343

3795

B16

Baker, Arthur Latham
Elliptic functions

Date Due			

University of Pittsburgh at Bradford

